

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/254021048>

# Recent advances in medicinal plant biotechnology

Article in *Indian Journal of Biotechnology* · January 2009

CITATIONS

59

READS

6,269

4 authors, including:



**Mohammad Yaseen Khan**

Nirma University

18 PUBLICATIONS 174 CITATIONS

SEE PROFILE



**Aliabbas Saleh**

National University of Ireland, Maynooth

8 PUBLICATIONS 92 CITATIONS

SEE PROFILE



**Shalini Rajkumar**

Nirma University

33 PUBLICATIONS 291 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Molecular Physiology of Plant Growth Promoting Rhizobacteria (PGPR) [View project](#)



herbal anticancer liposome [View project](#)

## Recent advances in medicinal plant biotechnology

Mohammad Yaseen Khan<sup>1</sup>, Saleh Aliabbas<sup>2</sup>, Vimal Kumar<sup>1</sup> and Shalini Rajkumar<sup>2\*</sup>

<sup>1</sup>Department of Phytopharmaceuticals and Natural Products, Institute of Pharmacy  
and

<sup>2</sup>Department of Biochemistry and Biotechnology, Institute of Science,  
Nirma University of Science and Technology, Sarkhej-Gandhinagar Highway, Ahmedabad 382 481, India

*Received 18 July 2007; revised 19 May 2008; accepted 10 August 2008*

Medicinal plants are the most important source of life saving drugs for the majority of the world's population. Plant secondary metabolites are economically important as drugs, fragrances, pigments, food additives and pesticides. The biotechnological tools are important to select, multiply, improve and analyze medicinal plants. *In-vitro* production of secondary metabolites in plant cell suspension cultures has been reported from various medicinal plants and bioreactors are the key step towards commercial production of secondary metabolites by plant biotechnology. Genetic transformation is a powerful tool for enhancing the productivity of novel secondary metabolites; especially by *Agrobacterium tumefaciens*. Combinatorial biosynthesis is another approach in the generation of novel natural products and for the production of rare and expensive natural products. DNA profiling techniques like DNA microarrays serve as suitable high throughput tools for the simultaneous analysis of multiple genes and analysis of gene expression that becomes necessary for providing clues about regulatory mechanism, biochemical pathways and broader cellular functions.

**Keywords:** Medicinal plants, biotechnology, combinatorial biosynthesis, DNA microarray, transgenic plants

### Introduction

It is estimated that 70-80% of people worldwide rely chiefly on traditional, largely herbal, medicines to meet their primary healthcare needs<sup>1</sup>. The global demand for herbal medicine is not only large, but growing<sup>1,2</sup>. Various technologies have been adopted for enhancing bioactive molecules in medicinal plants<sup>3</sup>. Biotechnological tools are important for the multiplication and genetic enhancement of the medicinal plants by adopting techniques such as *in vitro* regeneration and genetic transformation<sup>4</sup>. It could also be harnessed for the production of secondary metabolites using plants as bioreactors. Advances in tissue culture, combined with improvement in genetic engineering techniques specifically transformation technology, have opened new avenues for high volume production of pharmaceuticals, nutraceuticals and other beneficial substances. Recent advances in the molecular biology, enzymology and fermentation technology of plant cell cultures suggest that these systems may become a viable source of important secondary

metabolites. DNA manipulation is resulting in relatively large amounts of desired compounds produced by plants infected with an engineered virus, whereas transgenic plants can maintain constant levels of production of proteins without additional intervention. Large-scale use of plant tissue culture is found to be an attractive alternative approach to traditional methods of plantation as it offers a controlled supply of biochemicals independent of plant availability<sup>5,6</sup>. Impact of specific engineering-related factors on cell suspension cultures has also been detailed<sup>6</sup>. Current developments in tissue culture technology indicate that transcription factors are efficient new molecular tools for plant metabolic engineering to increase the production of valuable compounds<sup>7</sup>. The approach to combine genes from different microorganisms for the production of new and interesting metabolites is known as combinatorial biosynthesis, which has emerged as a new tool in the generation of novel natural products, as well as for the production of rare and expensive natural products. There are several pharmaceuticals on the market that are highly expensive, due to the fact that these compounds are only found in rare plants and often in extreme low concentrations. Combinatorial biosynthetic strategies

\*Author for correspondence:

Tel: 91-2717-241901-04 extn 627; Fax: 91-2717-241916  
shalini\_rjk@yahoo.com

are expected to yield interesting alternatives in the near future. Combinatorial biosynthesis has been utilised for important classes of natural products, including alkaloids (vinblastine, vincristine), terpenoids (artemisinin, paclitaxel) and flavonoids. Conventional strategies for expression profiling such as northern blot, reverse northern blot, reverse transcriptase-polymerase chain reaction (RT-PCR), nuclease protection, enzyme-linked immunosorbent assay (ELISA), western blot, *in situ* hybridization and immunohistochemistry are optimized for single gene analysis. Although, it is possible to modify at least some of these techniques for multiplexing, the procedure becomes increasingly technically cumbersome. For genome wide expression analysis it is necessary to develop technologies having high degree of automation, since in any living organism thousands of genes and their products function in a complicated and orchestrated way. DNA microarrays have been developed in response to the need for a high-throughput, efficient and comprehensive strategy that can simultaneously measure all the genes, or a large defined subset encoded by a genome<sup>8,9</sup>. DNA microarrays are being used to study the transcriptional profile in various physiological and pathological conditions, leading to mining of novel genes and molecular markers for diagnosis, prediction or prognosis of those specific states<sup>10</sup>. This paper reviews the recent achievements and advances in the medicinal plant biotechnology.

### Plant Tissue Culture

Plant tissue culture refers to growing and multiplication of cells, tissues and organs of plants on defined solid or liquid media under aseptic and controlled environment. Plant regeneration through somatic embryogenesis from stem, petiole and leaf explants of Indian chicory (*Cichorium intybus* L.) has been achieved<sup>11,12</sup>. The commercial technology is primarily based on micropropagation, in which rapid proliferation is achieved from tiny stem cuttings, axillary buds and to a limited extent from somatic embryos, cell clumps in suspension cultures and bioreactors. The cultured cells and tissues can take several pathways. The pathways that lead to the production of true-to-type plants in large numbers are the preferred ones for commercial multiplication. The process of micropropagation is usually divided into several stages such as pre-propagation, initiation of explants, subculture of explants for proliferation, shooting and rooting and hardening. These stages are

universally applicable in large-scale multiplication of plants. Micropropagation (*in vitro* propagation of axillary and/or adventitious buds as well as somatic embryos) is presently used as an advanced biotechnological system for the production of identical pathogen-free plants for agriculture and forestry<sup>13</sup>. Protocols for the cloning of some medicinal plants such as *Catharanthus roseus* (Apocynaceae), *Chlorophytum borivilianum* (Liliaceae), *Datura metel* (Solanaceae) and *Bacopa monnieri* (Scrophulariaceae) have been developed<sup>14</sup>. Moreover, *in vitro* flowering, *in vitro* fruiting and effective micropropagation protocols were studied in *Withania somnifera*, an antitumor medicinal plant using axillary buds implants<sup>15</sup>. A rapid micropropagation protocol was developed for *Hoslundia opposita* using nodal explants derived from mature trees<sup>16</sup>. The integrated approaches of plant culture systems will provide the basis for the future development of novel, safe, effective, and high-quality products for consumers.

Micropropagation by conventional techniques is typically a labour-intensive means of clonal propagation. Automation of micropropagation in bioreactors has been advanced by several authors as a possible way of reducing costs of micropropagation<sup>17-19</sup>. Bioreactors containing liquid media are used for large-scale growth of various tissues. The use of bioreactor for the micropropagation was first reported in 1981 for *Begonia*<sup>20</sup>. Since then it has proved applicable to many species including shoot, bulbs, microtubers, corms and somatic embryos<sup>21</sup>. Bioreactors are vessels designed for large-scale cell, tissue or organ culture in liquid media. Functionally, plant culture bioreactors can be divided into two broad types: those in which the cultures are immersed partially or temporarily in the medium and those in which the cultures are continuously submerged. Bioreactors provide more precise control of the plant growth gaseous exchange, illumination, medium agitation, temperature and pH than the conventional culture vessels. Bioreactor-based propagation of plants can increase the rate of multiplication and growth of cultures and reduce the space, energy and labour requirements in commercial micropropagation. Three main classes of culture systems in bioreactors can be distinguished: (1) those producing biomass (cells or organogenic or embryogenic propagules, shoots or roots as the final product); (2) those producing metabolites and enzymes; and (3) biotransformation of exogenously

added metabolites (which may be precursors in the pathway). Less labour-intensive clonal propagation at lower cost through the use of modified air-lift, bubble column, bioreactors (a balloon-type bubble bioreactor) together with temporary immersion systems for the propagation of shoots, bud-clusters and somatic embryos have been developed. The bioreactor system has been applied for embryogenic and organogenic cultures of several plant species<sup>22,23</sup>. Significant amount of sanguinarine was produced in cell suspension cultures of *Papaver somniferum* using bioreactors<sup>24</sup>. Ginseng root tissues, when cultured in a 20 tonne bioreactor, produced 500 mg/L/d of the saponin, which is considered as a very good yield<sup>25</sup>. Mass production of transformed *Panax ginseng* hairy roots in bioreactor has also been reported<sup>26</sup>. Much progress has been achieved in the recent past on optimization of these systems for the production and extraction of valuable medicinal plant ingredients such as ginsenosides and shikonin. Roots cultivated in bioreactors have been found to release medicinally active compounds, including the anticancer drug isolated from various *Taxus* species into the liquid media of the bioreactor, which may then be continuously extracted for pharmaceutical preparations<sup>27</sup>. Some of the recently produced bioactive secondary metabolites through plant tissue culture are presented in Table 1.

### Combinatorial Biosynthesis

The basic concept of combinatorial biosynthesis is to combine metabolic pathways in different organisms at the genetic level, where genes from different microorganisms are combined for the production of new and interesting plant secondary metabolites (Fig. 1)<sup>45</sup>. From pharmaceutical point of view, hydroxylations and glycosylations are considered to be particularly useful bioconversions<sup>38,39</sup>. They can yield new drugs and existing drugs can be improved as to increased activity and decreased toxicity. Recent achievements with the polyketide biosynthesis from microorganisms, especially in *Streptomyces*, prove the potential of combinatorial biosynthesis<sup>40-43</sup>. Podophyllotoxin and paclitaxel are clear examples of pharmaceuticals that can only be produced through the isolation from plants. With regard to the production of podophyllotoxin it has been shown that plant cell cultures of *Linum flavum* can be used to convert deoxypodophyllotoxin, a major lignan of *Anthriscus sylvestris* into 6-methoxypodophyllotoxin<sup>44</sup>. The combination of the product of one

Table 1—Recently produced bioactive secondary metabolites from plant tissue cultures

Plant name	Active ingredient	Culture type	References
<i>Cassia acutifolia</i>	Anthraquinones	Suspension	28
<i>Catharanthus roseus</i>	Catharanthine	Suspension	29
<i>Mentha arvensis</i>	Terpenoid	Shoot	30
<i>Nothapodytes foetida</i>	Camptothecin	Callus	31
<i>Podophyllum hexandrum</i>	Podophyllotoxin	Suspension	32
<i>Rhus javanica</i>	Gallotannins	Root	33
<i>Salvia fruticosa</i>	Rosmarinic acid	Callus & suspension	34
<i>Silybum marianum</i>	Flavonolignan	Root	35
<i>Taxus</i> species	Taxol	Suspension	36
<i>Withania somnifera</i>	Withaferin	Shoot	37

species and the enzymes of another species to yield a desired product forms a good example of combinatorial biosynthesis.

### Combinatorial Biosynthesis of Terpenoids

Terpenoids represent a large and important class of natural products with more than 30,000 different structures. From pharmaceutical point of view the sesquiterpenoids are of high relevance. In this group artemisinin, gossypol and zingiberene are of great medicinal and economic interest. Terpenoids are biosynthesized via mevalonate (MVA) pathway or deoxyxylulose phosphate (DOXP) pathway. The MVA pathway has recently been expressed in *E. coli* harbouring the DOXP pathway, which led to an efficient production of the terpenoids, amorpho-4,11-diene and taxadiene<sup>46-50</sup>.

### Artemisinin

Artemisinin is an antimalarial drug isolated from *Artemisia annua* (Asteraceae). The selection of plants yielded varieties containing 0.5-1.16% of artemisinin in the aerial parts based on dry weight<sup>51,52</sup>. Alternatives could be the production via transgenic plants<sup>53,54</sup> or engineering the biosynthetic pathway into less complex host cells. This implies that the full elucidation of the biosynthetic pathway is required. Although several biosynthetic pathways have been postulated, until now only the genes encoding the enzymes for the synthesis of the first specific




Fig. 1—Combinatorial biosynthesis

intermediate amorphadiene by amorphadiene synthase<sup>55,56</sup> and artemisinic acid by the cytochrome P450 enzyme, CYP71AV1<sup>57</sup> have been isolated and identified. The cDNA encoding amorphadiene synthase has been expressed in *E. coli* and characterized<sup>58</sup>. The recently discovered enzyme, CYP71AV1 was shown to catalyze the regioselective oxidation of amorphadiene into artemisinic alcohol. Besides this metabolic action, the enzyme also oxidizes the precursors artemisinic alcohol and artemisinic aldehyde yielding artemisinic acid<sup>58</sup>.

#### Paclitaxel

Paclitaxel, mostly described by the tradename Taxol, is a diterpenoid that can be found in the bark and needles of different species of *Taxus*. The biosynthesis of paclitaxel starts with the cyclization step from geranylgeranyl diphosphate (GGDP) to taxadiene. Most of the 19 known enzymatic steps in the biosynthesis are related to hydroxylation and other oxygenation reactions of the taxadiene skeleton<sup>59,60</sup>. Several genes from different *Taxus* species that are responsible for steps in the biosynthesis and building a basis for today's combinatorial biosynthesis in a heterologous microorganism have been isolated and

identified. Today, all the genes have been cloned into *E. coli* and activity screening confirmed the function of isolated enzymes<sup>61-74</sup>. The first intermediate, taxadiene can now be produced in *E. coli*. Co-expression of the taxadiene synthase from *Taxus brevifolia*<sup>75</sup> with a geranylgeranyl diphosphate synthase isolated from *Erwinia herbicola*<sup>76</sup>, isopentenyl diphosphate synthase from *Schizosaccharomyces pombe*<sup>77</sup>, and the endogenous deoxyxylulose 5-phosphate synthase from *E. coli* resulted in the production of 1.3 mg/L taxadiene in cell culture<sup>78</sup>.

#### Carotenoids

Combinatorial biosynthesis of carotenoids in microorganisms has also been described<sup>79</sup>. The yeast, *Candida utilis*, has been engineered for the production of lycopene,  $\beta$ -carotene, and astaxanthin<sup>80</sup>. The production of carotenoids in a host requires the biosynthesis of the intermediate GGDP. *E. coli* produces the C15 precursor FDP for endogenous terpenoid molecules. The extension of the prenyl chain to C20 has been performed by the expression of CrtE gene encoding geranylgeranyl diphosphate synthase from *Erwinia* sp.<sup>81</sup>. Prenyltransferase

catalyses the production of GGDP from FDP; the GGDP synthase encoding gene *gps* from *Archaeoglobus fulgidis* has been expressed as well. Expression of this gene is more efficient, because the enzyme catalyzes the three chain elongation reactions starting from C5 precursors to C20 molecule<sup>82</sup>.

#### Combinatorial Biosynthesis of Alkaloids

Combinatorial biosynthesis has been reported for the alkaloids, vincristine, vinblastine, ajmaline and morphine from plants and for rebeccamycin and staurosporine from *Streptomyces albus*<sup>83-85</sup>. The morphine biosynthesis consists of 17 steps in *Papaver somniferum*. In the biosynthesis, a key intermediate, (S)-norcoclaurine is biosynthesized by condensation of dopamine and 4-hydroxyphenylacetaldehyde (4-HPAA). The catalyzing enzyme (S)-norcoclaurine synthase has recently been identified from *Thalictrum flavum* and cloned in *E. coli*<sup>86,87</sup>. Further, key enzymatic steps towards (S)-reticuline include three NADPH oxidoreductases and cytochrome P450<sup>88,89</sup> and an acetyl-CoA dependent acetyltransferase<sup>90,91</sup>. Recently, the last step reducing codeine to morphine by codeinone reductase has been elucidated and the gene expressed in *E. coli* and insect cells<sup>92</sup>.

#### *Vinca* Alkaloids

Vinblastine and vincristine, the monoterpenoid indole alkaloids from *Catharanthus roseus*, are used as antineoplastic drugs. Because of the high importance and the extreme low yields from plants (3 mg kg<sup>-1</sup>) they could be considered as trace compounds<sup>93</sup>. *In vitro* application to improve alkaloid yield in *C. roseus* has been reviewed by many authors<sup>94</sup>. It is estimated that for the production of 3 kg of *Vinca* alkaloids, which is the annual need worldwide, around 300 tons of plant material has to be extracted. Production of *Vinca* alkaloids in plant cell cultures did not lead to a significant improvement and today it is accepted that biotechnological approaches in plant cell culturing may not provide an instant solution to this problem. The biosynthesis of vincristine and vinblastine is complex, from early phase starting from geraniol to strictosidine and for the late phase leading to the desired compounds. In the initial phase, tryptophan and secologanin as terpenoid precursors are condensed to form strictosidine as an important branching intermediate for other alkaloids. In this short part of the entire route seven enzymes and corresponding genes are involved, four of which have been cloned in *E. coli*<sup>95,96</sup>.

#### Genetic Transformation Technology and Production of Transgenic Plants

Genetic transformation technology has been proved to be a powerful tool for the production of plants with desired traits in many crops<sup>97,98</sup>. It promises to overcome some of the substantial agronomic and environmental problems that have not been solved using conventional plant breeding programmes<sup>99</sup>.

#### *Agrobacterium* and non-*Agrobacterium* Mediated Gene Transfer

Plant transformation mediated by *Agrobacterium tumefaciens*, a soil plant pathogenic bacterium, has become the most commonly used method for the introduction of foreign genes into plant cells and the subsequent regeneration of transgenic plants<sup>100</sup>. This soil bacterium possesses the natural ability to transform its host by delivering a well-defined DNA fragment, the transferred (T) DNA, of its tumour-inducing (Ti) plasmid into the host cell<sup>101,102</sup>. The rapid progress in the area of crop biotechnology is mainly because of the development of efficient regeneration and suitable *Agrobacterium*-mediated transformation protocols for different crop species<sup>103,104</sup>. Similar success could also be achieved in medicinal plants, which in turn could be used for the enhancement of secondary metabolite content. *Agrobacterium*-mediated genetic transformation of *Bacopa monniera* was standardized using the *A. tumefaciens* strain EHA105 that harboured the binary vector pBE2113 containing genes for glucuronidase (GUS) and neomycin phosphotransferase<sup>105</sup>. Transformation systems based on *A. tumefaciens* are well established for *Taxus* (yew)<sup>106</sup>, *Echinacea*<sup>107</sup>, *Scrophularia* (figwort)<sup>108</sup>, *Digitalis* (foxglove)<sup>109</sup>, *Thalictrum* (meadowrues)<sup>110</sup> and *Artemisia*<sup>111</sup>. An *A. tumefaciens*-mediated transformation system was developed for *Artemisia annua*. Analysis of artemisinin demonstrated that about 8-10 mg L<sup>-1</sup> DW of artemisinin was detected in the transgenic plants regenerated from five shoot lines, which is about 2-3 times higher than the control<sup>111</sup>. Thus, *Agrobacterium* transformation provides a method for routine genetic transformation of many important medicinal species. *Agrobacterium*-mediated gene transfer has the advantage for allowing stable integration of defined DNA into the plant genome that generally results in lower copy number, fewer rearrangements and more stability of expression over generations than other transfection methods<sup>112,113</sup>. Recent studies have shown that virus-based vectors can be efficiently used for high transient expression of foreign proteins in

transfected plants and that non-*Agrobacterium* bacterial species can be used for the production of transgenic plants, laying the foundation for alternative tools for future plant biotechnology. Non-*Agrobacterium* species like *Rhizobium* sp. NGR234, *Sinorhizobium meliloti* and *Mesorhizobium loti*, are capable of genetically transforming different plant tissues and plant species<sup>114</sup>.

#### Direct Gene Transfer

##### *Generation of Transgenic Medicinal Plants by Particle Bombardment*

Particle bombardment procedure was introduced in 1987, which involves the use of a modified shotgun to accelerate small (1-4  $\mu\text{m}$ ) diameter metal particles into plant cells at a velocity sufficient to penetrate the cell wall. There is no intrinsic limitation to the potential of particle bombardment since DNA is governed entirely by physical parameters. Different types of plant materials have been used as transformation targets including callus, cell suspension cultures and organised tissues such as immature embryos and meristems.

Cell suspension cultures were established from leaf explants of gentian (*Gentiana triflora*  $\times$  *G. scabra*) for generation of transgenic plants by particle bombardment<sup>115</sup>. Efficient transformation of the tropane alkaloid-producing medicinal plant, *Hyoscyamus muticus*, was also achieved by particle bombardment<sup>116</sup>. An efficient and stable transformation has been achieved in garlic plants (*Allium sativum*)<sup>117</sup>. The results indicate that biolistic transformation can lead to the transfer, expression and stable integration of a DNA fragment into chromosomal DNA. The relative simplicity of this system is a good recommendation for its future use in the production of genetically modified plants.

##### *Generation of Transgenic Medicinal Plants by Electroporation*

Electroporation uses brief pulses of high voltage electricity to induce the formation of transient pores in the membrane of the host cell. Such pores appear to act as passageways through which the naked DNA can enter the host cell. Exposure of cell suspension protoplasts of the woody medicinal plant, *Solatum dulcamara*, to a voltage of 250 to 1250 V  $\text{cm}^{-1}$  for three successive pulses, each of 10-50  $\mu\text{s}$  duration, stimulated growth of protoplast-derived tissues<sup>118</sup>. Such tissues exhibited increased morphogenesis and required a shorter period in culture to exhibit this effect than tissues from untreated protoplasts.

Regenerated shoots also rooted more readily and developed more prolific root systems than shoots from untreated protoplasts. These observations have important implications for plant genetic manipulation and may have application in the recovery and rooting of shoots from tissues of woody species, normally considered recalcitrant in culture.

##### *Generation of Transgenic Medicinal Plants by Chloroplast Transformation*

Stable transformation of the chloroplast by inserting foreign genes into the chloroplast genome - was first achieved in the single cell green alga, *Chlamydomonas reinhardtii*<sup>119</sup> in 1988, soon to be followed by tobacco plant, and more recently, *Arabidopsis thaliana*. More than 40 transgenes have been stably integrated and expressed using the tobacco chloroplast genome to confer desired agronomic traits or express high levels of vaccine antigens and biopharmaceuticals<sup>120</sup>. Leaf discs are bombarded with plasmid constructs containing a selectable antibiotic resistance marker physically linked to the gene of interest, flanked by DNA for inserting into the correct site of the chloroplast genome. The antibiotic resistance marker most frequently used is the *aadA* gene encoding resistance for spectinomycin and streptomycin, driven by the promoter of the chloroplast encoded 16S rRNA.

##### *Advantages of Chloroplast Transformation*

1. There is no risk of the transgene being transmitted through pollen to plants.
2. Transgenes integrated into the chloroplast genome show very high levels of expression (up to 40% of the soluble protein in the cell may be the recombinant protein).
3. The transgene expression is not affected by gene silencing.
4. Chloroplasts express bacterial genes better than nuclei.
5. High level of transgene expression ensures a high level of mortality of the pests<sup>121</sup>.

##### *Pathway Engineering*

Increasing the production of active phytochemical constituents is a well established target for genetic manipulation but presents some severe challenges. In particular, the metabolic pathways by which active compounds are biosynthesized are mostly poorly understood and relatively few genes for key enzymatic or regulatory steps have been isolated.

Nevertheless, there are examples of pathway engineering leading to improvements of potential value in the breeding of medicinal plants<sup>122,123</sup>. A recent article illustrating the challenges and opportunities of this approach describes a nine-fold enhancement in the production of a sedative compound, scopolamine, in hairy root cultures of *Hyoscyamus niger* (black henbane), brought about by simultaneously over expressing two genes encoding the rate limiting upstream and downstream biosynthetic enzymes<sup>124</sup>. Increase in the production of scopolamine in *A. belladonna*, from the naturally occurring chemical precursor hyoscyamine, by transformation with the enzyme hyoscyamine 6  $\beta$ -hydroxylase from *Hyoscyamus* is reported<sup>125</sup>. Preliminary progress has also been made towards engineering alkaloid production in *P. somniferum*<sup>126</sup>. A three-fold enhancement in the production of the putative antimalarial, anti-cancer agent, artemisinin, has been reported in transgenic *Artemisia* plants overexpressing farnesyl di phosphate synthase, the enzyme immediately preceding the first committed biosynthetic step<sup>127,128</sup>. New genomic approaches and efficient gene isolation methods applied to difficult secondary pathways in medicinal plant metabolism will undoubtedly expand the range and precision of manipulations via transgenesis, providing potentially superior material for the breeder.

#### Engineering Agronomic Traits in Medicinal Plants

There have been relatively few reports of biotechnology applied to abiotic stresses and other aspects of agronomic performance of medicinal plants, although the approach is considered to have great potential<sup>129</sup>. Just as resistance to herbicides, pests and diseases are the characters that have led the way in the introduction of transgenic crop species, so too have these characters been among the first targets for medicinal plant biotechnology. Transgenic *Atropa* plants resistant to the herbicides, bi-alaphosandglufosinate, has been described<sup>130</sup> and *Panax ginseng*, resistant to the herbicide, Basta, has been generated by transformation with the genes encoding enzyme phosphinothricin acetyl-transferase<sup>131</sup>. Resistant biotypes can also be a useful germplasm source for breeding. For instance, somatic hybridization, the fusion of somatic cells from tissue culture and atrazine selection, have been used to regenerate herbicide tolerant *Solanum nigrum* (blacknightshade)<sup>132</sup>. Transgenesis will assist in breeding varieties resistant to pathogens especially

those causing fungal diseases. *Panax quinquefolium* (American ginseng) transformed with either a chitinase or thaumatin-like antifungal genes have been regenerated successfully<sup>133, 134</sup>.

#### Bioanalytics and Metabolomics

##### Latest Hyphenated Techniques of Spectroscopic Analysis

The coupling of high performance liquid chromatography with nuclear magnetic resonance spectroscopy (LC-NMR) is one of the most powerful methods for the separation and structural elucidation of unknown compounds in mixtures. LC-NMR thus represents a potentially interesting complementary technique to LC-UV-MS in phytochemical analysis for the detailed on-line structural analysis of natural products. *Ligusticum chuanxiong* Hort., a plant that is frequently used in traditional Chinese medicine, has been studied using HPLC-coupled spectroscopic techniques such as HPLC-UV, HPLC-MS as well as HPLC-NMR<sup>135,136</sup>.

##### Genomic Profiling of Medicinal Plants by DNA Microarray Technique

DNA microarray was developed in response to the need for a high-throughput, efficient and comprehensive strategy that can simultaneously measure all the genes or a large defined subset, encoded by a genome<sup>137,138</sup>. Several methodologies including differential display PCR, northern blots<sup>139</sup>, quantitative PCR, serial analysis of gene expression (SAGE)<sup>140,141</sup> and TIGR Orthologous Gene Alignments (TOGA)<sup>142, 143</sup> are used along side microarrays as research tools. DNA microarray is an orderly arrangement of thousands of oligonucleotides or identified sequenced genes printed on an impermeable solid support, usually glass, silicon chips or nylon membrane. Novel technologies such as SAGE and DNA microarray allow rapid and detailed analysis of thousands of transcripts, providing a revolutionary approach to the investigation of gene expression. There are three main applications of DNA microarray:

- (i) in pharmacodynamics for discovery of new diagnostic and prognostic indicators and biomarkers of therapeutic response; elucidation of molecular mechanism of action of an herb, its formulations or its phytochemical components and identification and validation of new molecular targets for herbal drug development.
- (ii) in pharmacogenomics for prediction of potential side-effects of the herbal drug during preclinical activity and safety studies;



identification of genes involved in conferring drug sensitivity or resistance and prediction of patients most likely to benefit from the drug and use in general pharmacogenomic studies.

- (iii) in pharmacognosy for correct botanical identification and authentication of crude plant materials as part of standardization and quality control.

The identification of genes modulated by compounds isolated from *Centella asiatica* with the help of gene microarray provides the basis for a molecular understanding of *Centella's* bioactivity, and opportunities for the quantitative correlation of this activity with clinical effectiveness at molecular level<sup>144</sup>. Similarly, the antiproliferative activity of *Coptidis rhizoma*, a medicinal herb and its major component berberine was investigated in human pancreatic cancer cell lines and it was possible to identify common and distinct genes related to anti-proliferative activities<sup>145</sup>. High-density oligonucleotide microarrays have been used for pioneering studies on the multiple gene expression effects exhibited by *Ginkgo biloba* leaf extract EGb 761<sup>146</sup> and various studies on diets supplemented with *G. biloba* extract that have notable neuromodulatory effects *in vivo* and illustrates the utility of genome-wide expression monitoring to investigate the biological actions of complex extracts<sup>147</sup>.

cDNA microarray analyses have shown that exposure of human breast cancer cells to a *Ginkgo* extract altered the expression of genes that are involved in the regulation of cell proliferation, cell differentiation or apoptosis, and that exposure of human bladder cancer cells to a *Ginkgo* extract produced an adaptive transcriptional response that augments antioxidant status and inhibits DNA damage<sup>148</sup>. Recent studies have highlighted the concurrent use of herbs which may mimic, magnify, or oppose the effect of drugs<sup>149,150</sup>. DNA microarray can be used for studying herb–drug interactions, and the mechanisms underlying these interactions.

#### DNA Microarray in Pharmacognosy

Use of an authentic herbal material is the first step of ensuring quality, safety and efficacy of herbal medicines. DNA polymorphism-based assays have been developed for the identification of herbal medicines<sup>151,152</sup>. Recently, microarrays have been applied for the DNA sequence-based identification of medicinal plants<sup>153,154</sup>. This includes quality control and standardization of the herbal drugs, identification

and validation of new targets, the profiling of on-target and off-target effects during the optimization of new therapeutic agents, understanding molecular mechanisms of action, structure–activity relationships<sup>155</sup> and the prediction of side-effects, and the discovery of diagnostic, prognostic, and pharmacodynamic biomarkers. Oligonucleotide probes specific for polymorphisms in the D2 and D3 regions of 26S rDNA gene of several *Fritillaria* species were designed and printed on the poly-lysine coated slides to prepare a DNA chip. Differentiation of the various *Fritillaria* species was accomplished based on hybridization of fluorescent labeled PCR products with the DNA chip. The results demonstrated the reliability of using DNA chips to identify different species of *Fritillaria*, and that the DNA chip technology can provide a rapid, high throughput tool for genotyping and plant species authentication<sup>156</sup>. Similarly, using fluorescence-labeled ITS2 sequences as probes, distinctive signals were obtained for the five medicinal species of *Dendrobium* listed in the Chinese Pharmacopoeia. It established that microarray was able to detect the presence of *D. nobile* in a Chinese medicinal formulation containing nine herbal components<sup>157</sup>. Phytomics, a technology platform for characterization of herbal compositions wherein, herbal bio response arrays (HBR Arrays) are used to determine bioactive constituents and biological activities of an herbal composition has recently been developed and patented<sup>158</sup>.

cDNA-amplified fragment length polymorphism (AFLP) has been the functional-genomics tool of choice to study gene expression profiles related to the biosynthesis of secondary metabolites. cDNA-AFLP has the advantage as it does not require prior genomic data<sup>159,160</sup>. For most medicinal plants either limited or no information on genomic sequences is available, or cDNA libraries that could be used as a template for microarrays do not exist. cDNA-AFLP is done by cDNA synthesis from the tissue of choice and selective amplification. Genome wide profiling of transcript is done using softwares such as AFLP Quantarpro (Keygene, Wageningen, The Netherlands).

#### Marker Assisted Breeding

Marker assisted breeding involves the use of DNA markers linked with DNA sequences of interest. Inheritance pattern of sequences/traits can be confirmed even prior to expression using RFLP,

RAPD, AFLP and minisatellite markers<sup>161,162</sup>. A high content artemisinin producing plant variety “CIM-Arogya” was developed at CIMAP, Lucknow through marker assisted breeding. Selection of genotype with increased biomass led to selection of higher artemisinin yielding variety. An increase in artemisinin from 0.15% to 1.16% on dry weight basis was recorded in this variety<sup>163</sup>. Thus, marker assisted breeding could also be used in developing high secondary metabolite yielding plants.

#### Expressed Sequence Tags (ESTs) of Medicinal Plants

Monoterpene indole alkaloid (MIA) pathway genes were identified from random sequencing of *C. roseus* cDNA library which revealed 3655 unique ESTs, composed of 1142 clusters and 2513 singletons. Several novel MIA pathway candidate genes were identified by the cloning and functional characterization of loganic acid O-methyltransferase involved in secologanin biosynthesis. Biochemical pathways such as triterpene biosynthesis were also identified and its metabolite analysis revealed localization of oleanane-type triterpenes exclusively to the cuticular wax layer. The results illuminated, biochemical specialization of *Catharanthus* leaf epidermis for the production of multiple classes of metabolites<sup>164</sup>. Also, jasmonate-induced changes on the transcript and alkaloid profiles of tobacco BY-2 and *C. roseus* cell cultures have been monitored through the similar approach<sup>165,166</sup>. ESTs corresponding to 40 enzymes involved in the conversion of sucrose to sanguinarine were identified from elicitor induced cell culture of *Papaver somniferum*. Substantial increase in the level of RNA was observed in case of elicited cell culture as compared to control and the identified metabolites were sanguinarine, dihydrosanguinarine, methoxylated derivatives dihydrochelirubine and chelirubine, and the alkaloid pathway intermediates N-methylcoclaurine, N-methylstylopine, and protopine<sup>167</sup>. Similarly, cDNA library of *Artimisia annua* glandular trichome revealed the presence of many ESTs involved in isoprenoid biosynthesis such as enzymes from the methylerythritol phosphate pathway and the mevalonate pathway, amorpha-4,11-diene synthase and other sesquiterpene synthases, monoterpene synthases and two cDNAs showing high similarity to germacrene A synthases<sup>168</sup>. An inventory of hundreds of genes, potentially involved not only in alkaloid biosynthesis but also possibly in plant secondary metabolism in general, has been built.

Thereafter, large-scale functional analysis of genes from this inventory, potentially involved in plant secondary metabolism, was performed. This includes isolation, introduction and functional analysis of full-length open reading frames (FLORFs) in transgenic plant cells. Tools to improve and speed-up functional analysis of candidate genes in transgenic plant cells such as medium-throughput strategies for isolation of FLORFs, super-transformation of plant cells with reporter gene constructs, transient protoplast expression assays and microarray facilities, have been designed and their use validated.

#### Conclusion

The improved *in vitro* plant cell culture systems have the potential for commercial exploitation of secondary metabolites. Micropropagation, combined with *Agrobacterium* transformation, provides a method for routine genetic transformation of many important medicinal species. The production of secondary metabolites could be enhanced using bioreactors and has a tremendous potential for the large-scale synthesis of therapeutically active compounds in medicinal plants. During the past decade, remarkable progress in plant genetic-transformation technology has been witnessed. This rapid progress has resulted in constant flow of new and improved transformation protocols for many medicinal plant species. Genetic transformation may provide increased and efficient system for *in vitro* production of secondary metabolites. This review also highlights the possibilities for the use of bioconversion and combinatorial biosynthesis strategies for the production and development of plant natural products at different levels in biosynthetic pathways. There are several benefits of using microorganisms instead of plants or plant cell cultures, including their fast replication, low costs of cultivation, and the possibility for bioprocessing on an industrial scale. Microarray analysis of gene expression could be useful for elucidating the molecular mechanisms and networks underlying the complex pharmacological function of herbal extracts and mixtures. DNA microarrays have the potential for applications in different phases of herbal drug discovery and development. The tools described in this review would certainly be of increasing importance in the field of medicinal plant biotechnology research in near future.

#### References

- 1 Farnsworth N R & Soejarto D D, Global importance of medicinal plants, in *Conservation of medicinal plants*, edited

- by O Akerele, V Heywood & H Synge (Cambridge University Press, Cambridge) 1991, 25-51.
- 2 Srivastava R, Studying the information needs of medicinal plant stakeholders, *Traffic Dispatches*, 15 (2000) 5.
  - 3 Abdin M Z, Enhancing bioactive molecules in medicinal plants, in *Natural products—Essential resources for human survival*, edited by Y Zhu, B Tan, B Bay & C Liu (World Scientific Publishing Co. Pvt. Ltd., Singapore) 2007, 45-57.
  - 4 Abdin M Z & Kamaluddin, Improving quality of medicinal herbs through physio-chemical and molecular approaches, in *Traditional systems of medicine*, edited by M Z Abdin & Y P Abrol (Narosa Publishing House Pvt. Ltd., India) 2006, 30-39.
  - 5 Sajc L, Grubisic D & Vunjak-Novakovic G, Bioreactors for plant engineering: An outlook for further research, *Biochem Eng J*, 4 (2000) 89-99.
  - 6 Kieran P M, MacLoughlin P F & Malone D M, Plant cell suspension cultures: Some engineering considerations, *J Biotechnol*, 59 (1997) 39-52.
  - 7 Gantet, P & Memelink J, Transcription factors: Tools to engineer the production of pharmacologically active plant metabolites, *Trends Pharmacol Sci*, 23 (2002) 563-569.
  - 8 Schena M, Shalon D, Davis R W & Brown P O, Quantitative monitoring of gene expression patterns with a complementary DNA microarray, *Science*, 270 (1995) 467-470.
  - 9 Schena M, Shalon D, Heller R, Chai A, Brown P O *et al*, Parallel human genome analysis: Microarray-based expression monitoring of 1000 genes, *Proc Natl Acad Sci USA*, 93 (1996) 10614-10619.
  - 10 Gebauer M, Microarray applications: Emerging technologies and perspectives, *Drug Discov Today*, 9 (2004) 915-917.
  - 11 Abdin M Z & Ilah A, Plant regeneration and somatic embryogenesis from stem and petiole explants of Indian chicory (*Cichorium intybus* L.), *Indian J Biotechnol*, 6 (2007) 250-255.
  - 12 Rehman R U, Israr M, Srivastava P S, Bansal K C & Abdin M Z, *In vitro* regeneration of witloof chicory (*Cichorium intybus* L.) from leaf explants and accumulation of esculin, *In Vitro Cell Dev Biol*, 39 (2003) 142-146.
  - 13 George E F, *Plant propagation by tissue culture*, Part 2. (Exegetic Ltd, Edington Wesbury, UK) 1993.
  - 14 Mousumi D, Malik C P & Bisen P S, Micropropagation: A tool for the production of high quality plant-based medicines, *Curr Pharm Biotechnol*, 7 (2006) 33-49.
  - 15 Saritha K V & Naidu C V, *In vitro* flowering of *Withania somnifera* Dunal.—An important antitumor medicinal plant, *Plant Sci*, 172 (2007) 847-851.
  - 16 Prakash S & Van Staden J, Micropropagation of *Hoslundia opposita* Vahl—A valuable medicinal plant, *S Afr J Bot*, 73 (2007) 60-63.
  - 17 Preil W, Application of bioreactors in plant propagation. in *Micropropagation: Technology and application*, edited by P C Debergh and R H Zimmerman (Kluwer Academic Publ., Dordrecht, The Netherlands) 1991, 425-455.
  - 18 Takayama S & Akita M, The types of bioreactors used for shoots and embryos, *Plant Cell Tissue Organ Cult*, 39 (1994) 147-156.
  - 19 Leathers R R, Smith M A L & Aitken-Christie J, Automation of the bioreactor process for mass propagation and secondary metabolism, in *Automation and environment control in plant tissue culture*, edited by J Aitken-Christie, T Kozai & M A L Smith (Kluwer Academic Publ., Dordrecht, The Netherlands) 1995, 87-214.
  - 20 Takayama S & Misawa M, Mass propagation of *Begonia hiemalis* plantlets by shake culture, *Plant Cell Physiol*, 22 (1981) 461-467.
  - 21 Paek K Y, Hahn E J & Son S H, Application of bioreactors for large-scale micropropagation systems of plants, *In vitro Cell Dev Biol Plant*, 37 (1981) 149-157.
  - 22 Levin R, Gaba V, Tal B, Hirsch S, De Nola D *et al*, Automated plant tissue culture for mass propagation, *Biotechnology*, 6 (1988) 1035-1040.
  - 23 Preil W, Florek P, Wix U & Beck A, Towards mass propagation by use of bioreactors, *Acta Hort*, 226 (1988) 99-105.
  - 24 Park J M & Yoon S Y, Production of sanguinarine by suspension culture of *Papaver somniferum* in bioreactors, *J Ferment Bioeng*, 74 (1992) 292-296.
  - 25 Charlwood B V & Charlwood K A, Terpenoid production in plant cell cultures, in *Ecological chemistry and biochemistry of plant terpenoids*, edited by J B Harborne & F A Tomas-Barberan (Clarendon Press, Oxford) 1991, 95-132.
  - 26 Jeong G T, Park D H, Hwang B, Park K, Kim S W *et al*, Studies on mass production of transformed *Panax ginseng* hairy roots in bioreactor, *Appl Biochem Biotechnol*, 98 (2002) 1115-1127.
  - 27 Wink M, Alfermann A W, Franke R, Wetterauer B, Distl M *et al*, Sustainable bioproduction of phytochemicals by plant *in vitro* cultures: Anticancer agents, *Plant Genet Res*, 3 (2005) 90-100.
  - 28 Nazif N M, Rady M R & Seif E1-Nasr M M, Stimulation of anthraquinone production in suspension cultures of *Cassia acutifolia* by salt stress, *Fitoterapia*, 71 (2000) 34-40.
  - 29 Zhao J, Zhu W & Hu Q, Enhanced catharanthine production in *Catharanthus roseus* cell cultures by combined elicitor treatment in shake flasks and bioreactors, *Enzyme Microb Technol*, 28 (2001) 673-681.
  - 30 Phatak S V & Heble M R, Organogenesis and terpenoid synthesis in *Mentha arvensis*. *Fitoterapia*, 73 (2002) 32-39.
  - 31 Thengane S R, Kulkarni D K, Shrikhande V A, Joshi S P, Sonawane K B *et al*, Influence of medium composition on callus induction and camptothecin(s) accumulation in *Nothapodytes foetida*, *Plant Cell Tissue Organ Cult*, 72 (2003) 247-251.
  - 32 Chattopadhyay S, Srivastava A K, Bhojwani S S & Bisaria V S, Production of podophyllotoxin by plant cell cultures of *Podophyllum hexandrum* in bioreactor, *J Biosci Bioeng*, 93 (2002) 215-220.
  - 33 Taniguchi S K, Yazaki R, Yabuuchi K & Kawakami H, Ito T *et al*, Galloylglucosides and riccionidin A in *Rhus javanica* adventitious root cultures, *Phytochemistry*, 53 (2000) 357-363.
  - 34 Karam N S, Jawad F M, Arikat N S & Shibli R A, Growth and rosmarinic acid accumulation in callus, cell suspension, and root cultures of wild *Salvia fruticosa*, *Plant Cell Tissue Organ Cult*, 73 (2003) 117-121.
  - 35 Alikaridis F D, Papadakis K, Pantelia & Kephelas T, Flavonolignan production from *Silybum marianum* transformed and untransformed root cultures, *Fitoterapia*, 71 (2000) 379-384.

- 36 Wu J, Wang C & Mei X, Stimulation of taxol production and excretion in *Taxus* spp cell cultures by rare earth chemical lanthanum, *J Biotechnol*, 85 (2001) 67-73.
- 37 Ray S and Jha S, Production of withaferin A in shoot cultures of *Withania somnifera*. *Planta Med*, 67 (2001) 432-436.
- 38 Knaeblein J, Biopharmaceuticals expressed in plants, in *Parmaceutical biotechnology - Drug discovery and clinical applications*, edited by O Kayser & R H Müller (Wiley-VCH, Weinheim) 2004, 34-56.
- 39 Ma J K, Barros E, Bock R, Christou P, Dale P J *et al*, Molecular farming for new drugs and vaccines. Current perspectives on the production of pharmaceuticals in transgenic plants, *EMBO Rep*, 6 (2005) 593-599.
- 40 Hranueli D, Cullum J, Basrak B, Goldstein P & Long P F, Plasticity of the streptomyces genome-evolution and engineering of new antibiotics, *Curr Med Chem*, 12 (2005) 1697-1704.
- 41 Moore B S, Kalaitzis J A & Xiang L, Exploiting marine actinomycete biosynthetic pathways for drug discovery, *Antonie Van Leeuwenhoek*, 87 (2005) 49-57.
- 42 Weber T, Welzel K, Pelzer S, Vente A & Wohlleben W, Exploiting the genetic potential of polyketide producing streptomycetes, *J Biotechnol*, 106 (2003) 221-232.
- 43 Pfeifer B A & Khosla C, Biosynthesis of polyketides in heterologous hosts, *Microbiol Mol Biol Rev*, 65 (2001) 106-118.
- 44 Koulman A, Beekman A C, Pras N & Quax W J, The bioconversion process of deoxypodophyllotoxin with *Linum flavum* cell cultures, *Planta Med*, 69 (2003) 739-744.
- 45 Michels P C, Khmelnitsky Y L, Dordick J S & Clark D S, Combinatorial biocatalysis: A natural approach to drug discovery, *Trends Biotechnol*, 16 (1998) 210-215.
- 46 Lange B M, Rujan T, Martin W & Croteau R, Isoprenoid biosynthesis: The evolution of two ancient and distinct pathways across genomes, *Proc Natl Acad Sci USA*, 97 (2000) 13172-13177.
- 47 Eisenreich W, Bacher A, Arigoni D & Rohdich F, Biosynthesis of isoprenoids via non-mevalonate pathway, *Cell Mol Life Sci*, 61 (2004) 1401-1426.
- 48 Rohdich F, Kis K, Bacher A & Eisenreich W, The non-mevalonate pathway of isoprenoids: Genes, enzymes and intermediates, *Curr Opin Chem Biol*, 5 (2001) 535-540.
- 49 Martin V J, Pitera D J, Withers S T, Newman J D & Keasling J D, Engineering a mevalonate pathway in *Escherichia coli* for production of terpenoids, *Nat Biotechnol*, 21 (2003) 796-802.
- 50 Huang Q, Roessner C A, Croteau R & Scott A I, Engineering *Escherichia coli* for the synthesis of taxadiene, a key intermediate in the biosynthesis of taxol, *Bioorg Med Chem*, 9 (2001) 2237-2242.
- 51 Laughlin J C, Agricultural production of artemisinin-A review, *Trans R Soc Trop Med Hyg*, 88 (1994) S21-S22.
- 52 Wallaart T E, Pras N, Beekman A C & Quax W J, Seasonal variation of artemisinin and its biosynthetic precursors in plants of *Artemisia annua* of different geographical origin: Proof for the existence of chemotypes, *Planta Med*, 66 (2000) 57-62.
- 53 Abdin M Z, Israr M, Rehman R U & Jain S K, Artemisinin, a novel antimalarial drug: Biochemical and molecular approaches for enhanced production, *Planta Med*, 69 (2003) 289-299.
- 54 Abdin M Z, Israr M, Kumar P A & Jain S K, Molecular approaches to enhance artemisinin content in *Artemisia annua* L., in *Recent progress in medicinal plants: Biotechnology and genetic engineering*, vol IV, edited by J N Govil, P A Kumar & V K Singh (SciTech Publishing, Raleigh, NC USA) 2002, 145-162.
- 55 Wallaart T E, Bouwmeester H J, Hille J, Poppinga L & Majiers N C, Amorpha-4,11-diene synthase: Cloning and functional expression of a key enzyme in the biosynthetic pathway of the novel antimalarial drug artemisinin, *Planta*, 212 (2001) 460-465.
- 56 Mercke P, Bengtsson M, Bouwmeester H J, Posthumus M A & Brodelius P E, Molecular cloning, expression and characterization of amorpha-4,11- diene synthase, a key enzyme of artemisinin biosynthesis in *Artemisia annua* L., *Arch Biochem Biophys*, 381 (2000) 173-180.
- 57 Teoh K H, Polichuk D R, Reed D W, Nowak G & Covello P S, *Artemisia annua* L. (Asteraceae) trichome-specific cDNAs reveal CYP71AV1, a cytochrome P450 with a key role in the biosynthesis of the antimalarial sesquiterpene lactone artemisinin, *FEBS Lett*, 580 (2006) 1411-1416.
- 58 Picaud S, Olofsson L, Brodelius M & Brodelius P E, Expression, purification, and characterization of recombinant amorpha-4,11-diene synthase from *Artemisia annua* L., *Arch Biochem Biophys*, 436 (2005) 215-226.
- 59 Walker K & Croteau R, Taxol biosynthetic genes, *Phytochemistry*, 58 (2001) 1-7.
- 60 Jennewein S & Croteau R, Taxol: biosynthesis, molecular genetics and biotechnological applications, *Appl Microbiol Biotechnol*, 57 (2001) 13-19.
- 61 Ketchum R E, Rithner C D, Qiu D, Kim Y S, Williams R M & Croteau R B, *Taxus* metabolomics: Methyl jasmonate preferentially induces production of taxoids oxygenated at C-13 in *Taxus x media* cell cultures, *Phytochemistry*, 62 (2003) 901-909.
- 62 Long R M & Croteau R, Preliminary assessment of the C13-side chain 2'-hydroxylase involved in taxol biosynthesis, *Biochem Biophys Res Commun*, 338 (2005) 410-417.
- 63 Jennewein S, Park H, Dejong J M, Long R M, Bollon A P & Croteau R B, Coexpression in yeast of *Taxus* cytochrome P450 reductase with cytochrome P450 oxygenases involved in taxol biosynthesis, *Biotechnol Bioeng*, 89 (2005) 588-598.
- 64 Walker K D, Klettke K, Akiyama T & Croteau R, Cloning, heterologous expression, and characterization of a phenylalanine aminomutase involved in taxol biosynthesis. *J Biol Chem*, 279 (2004) 53947-53954.
- 65 Jennewein S, Wildung M R, Chau M, Walker K & Croteau R, Random sequencing of an induced *Taxus* cell cDNA library for identification of clones involved in taxol biosynthesis, *Proc Natl Acad Sci USA*, 101 (2004) 9149-9154.
- 66 Chau M & Croteau R, Molecular cloning and characterization of a cytochrome P450 taxoid 2a-hydroxylase involved in taxol biosynthesis, *Arch Biochem Biophys*, 42 (2004) 748-757.
- 67 Jennewein S, Long R M, Williams R M & Croteau R, Cytochrome P450 taxadiene 5a-hydroxylase, a mechanistically unusual monooxygenase catalyzing the first oxygenation step of taxol biosynthesis, *Chem Biol*, 11 (2004) 379-387.
- 68 Jennewein S, Rithner C D, Williams R M & Croteau R, Taxoid metabolism: Taxoid 14β-hydroxylase is a cytochrome P450-dependent monooxygenase, *Arch Biochem Biophys*, 413 (2003) 262-270.

- 69 Walker K, Fujisaki S, Long R & Croteau R, Molecular cloning and heterologous expression of the C-13 phenylpropanoid side chain-CoA acyltransferase that functions in taxol biosynthesis, *Proc Natl Acad Sci USA*, 99 (2002) 12715-12720.
- 70 Walker K, Long R & Croteau R, The final acylation step in taxol biosynthesis: Cloning of the taxoid C13-side-chain N-benzoyltransferase from *Taxus*, *Proc Natl Acad Sci USA*, 99 (2002) 9166-9171.
- 71 Jennewein S, Rithner C D, Williams R M & Croteau R B, Taxol biosynthesis: Taxane 13 $\alpha$ -hydroxylase is a cytochrome P450- dependent monooxygenase, *Proc Natl Acad Sci USA*, 98 (2001) 13595-13600.
- 72 Walker K & Croteau R, Taxol biosynthesis: Molecular cloning of a benzoyl-CoA:taxane 2 $\alpha$ -O-benzoyltransferase cDNA from *Taxus* and functional expression in *Escherichia coli*, *Proc Natl Acad Sci USA*, 97 (2000) 13591-13596.
- 73 Walker K, Schoendorf A & Croteau R. Molecular cloning of a taxa-4(20),11(12)-dien-5 $\alpha$ -ol-O-acetyl transferase cDNA from *Taxus* and functional expression in *Escherichia coli*, *Arch Biochem Biophys*, 374 (2000) 371-380.
- 74 Walker K & Croteau R, Molecular cloning of a 10-deacetylbaicatin III-10-O-acetyl transferase cDNA from *Taxus* and functional expression in *Escherichia coli*, *Proc Natl Acad Sci USA*, 97 (2000) 583-587.
- 75 Wildung M R & Croteau R, A cDNA clone for taxadiene synthase, the diterpene cyclase that catalyzes the committed step of taxol biosynthesis, *J Biol Chem*, 271 (1996) 9201-9204.
- 76 Math S K, Hearst J E & Poulter C D, The crtE gene in *Erwinia herbicola* encodes geranylgeranyl diphosphate synthase, *Proc Natl Acad Sci USA*, 89 (1992) 6761-6764.
- 77 Hahn F M & Poulter C D, Isolation of *Schizosaccharomyces pombe* isopentenyl diphosphate isomerase cDNA clones by complementation and synthesis of the enzyme in *Escherichia coli*, *J Biol Chem*, 270 (1995) 11298-11303.
- 78 Huang Q, Roessner C A, Croteau R, Scott A I, Engineering *Escherichia coli* for the synthesis of taxadiene, a key intermediate in the biosynthesis of taxol, *Bioorg Med Chem*, 9 (2001) 2237-2242.
- 79 Sandmann G, Genetic manipulation of carotenoid biosynthesis: Strategies, problems and achievements, *Trends Plant Sci*, 6 (2001) 14-17.
- 80 Miura Y, Kondo K, Shimada H, Saito T, Nakamura K & Misawa N. Production of lycopene by the food yeast, *Candida utilis* that does not naturally synthesize carotenoid, *Biotechnol Bioeng*, 58 (1998) 306-308.
- 81 Misawa N, Nakagawa M, Kobayashi K, Yamano S, Izawa Y, Nakamura K & Harashima K, Elucidation of the *Erwinia uredovora* carotenoid biosynthetic pathway by functional analysis of gene products expressed in *Escherichia coli*, *J Bacteriol*, 172 (1990) 6704-6712.
- 82 Wang C W, Oh M K & Liao J C, Engineered isoprenoid pathway enhances astaxanthin production in *Escherichia coli*, *Biotechnol Bioeng*, 62 (1999) 235-241.
- 83 Warzecha H, Gerasimenko I, Kutchan T M & Stockigt J, Molecular cloning and functional bacterial expression of a plant glucosidase specifically involved in alkaloid biosynthesis, *Phytochemistry*, 54 (2000) 657-666.
- 84 Sanchez C, Mendez C & Salas J A, Engineering biosynthetic pathways to generate antitumor indolocarbazole derivatives, *J Ind Microbiol Biotechnol*, 33 (2006) 560-568.
- 85 Sanchez C, Zhu L, Brana A F, Salas A P, Rohr J *et al*, Combinatorial biosynthesis of antitumor indolocarbazole compounds, *Proc Natl Acad Sci USA*, 102 (2005) 461-466.
- 86 Samanani N, Liscombe D K & Facchini P J, Molecular cloning and characterization of norcoclaurine synthase, an enzyme catalyzing the first committed step in benzyloquinoline alkaloid biosynthesis, *Plant J*, 40 (2004) 302-313.
- 87 Samanani N & Facchini P J, Purification and characterization of norcoclaurine synthase. The first committed enzyme in benzyloquinoline alkaloid biosynthesis in plants, *J Biol Chem*, 277 (2002) 33878-33883.
- 88 Gerardy R & Zenk M H, Formation of salutaridine from (R)-reticuline by a membrane bound cytochrome P450 enzyme from *Papaver somniferum*, *Phytochemistry*, 32 (1993) 79-86.
- 89 Huang F C & Kutchan T M, Distribution of morphinan and benzo[c]phenanthridine alkaloid gene transcript accumulation in *Papaver somniferum*, *Phytochemistry*, 53 (2000) 555-564.
- 90 Lenz R & Zenk M H, Acetyl coenzyme A: Salutaridinol-7-O-acetyltransferase from *Papaver somniferum* plant cell cultures. The enzyme catalyzing the formation of thebaine in morphine biosynthesis, *J Biol Chem*, 270 (1995) 31091-31096.
- 91 Grothe T, Lenz R & Kutchan T M, Molecular characterization of the salutaridinol 7-O-acetyltransferase involved in morphine biosynthesis in opium poppy *Papaver somniferum*, *J Biol Chem*, 276 (2001) 30717-30723.
- 92 Zhang Q, Rich J O, Cotterill I C, Pantaleone D P & Michels P C, 14-Hydroxylation of opiates: Catalytic direct autoxidation of codeinone to 14-hydroxycodeinone, *J Am Chem Soc*, 127 (2005) 7286-7287.
- 93 Van der Heijden R, Jacobs D I, Snoeijs W, Hallared D & Verpoorte R, The *Catharanthus* alkaloids: Pharmacognosy and biotechnology, *Curr Med Chem*, 11 (2004) 607-628.
- 94 Mujib A, Ilah A, Gandotra N & Abidin M Z, *In vitro* application to improve alkaloid yield in *Catharanthus roseus*, in *Recent progress in medicinal plants: Biotechnology and genetic engineering*, edited by J N Govil, P A Kumar & V K Singh (Sci. Tech. Pub., USA) 2002, 415-440.
- 95 Verpoorte R, Van der Heijden R, Schripsema J, Hoge J H C & Ten Hoopen H J G, Plant-cell biotechnology for the production of alkaloids- Present status and prospects, *J Nat Prod*, 56 (1993) 186-207.
- 96 Kutchan T M, Bock A & Dittrich H, Heterologous expression of the plant proteins strictosidine synthase and berberine bridge enzyme in insect cell culture, *Phytochemistry*, 35 (1994) 353-360.
- 97 Sawahel, W. *Plant genetic transformation technology* (Daya Publishing House, India) 1997.
- 98 Sawahel W & Cove D, Gene transfer strategies in plants, *Biotechnol Adv*, 10 (1992) 393-412.
- 99 Gasser C & Fraley R, Genetic engineering plants for crop improvement, *Science*, 244 (1989) 1293-1299.
- 100 Riva G A, Gonzalez-Cabrera J, Vazquez-Padrón R & Ayra-Pardo C, *Agrobacterium tumefaciens*: A natural tool for plant transformation, *J Biotechnol*, 3 15 (1998) 118-133.
- 101 Tzfira T, Li J, Lacroix B & Citovsky V, *Agrobacterium* T-DNA integration: Molecules and models, *Trends Genet*, 20 (2004) 375-383.
- 102 Gelvin S B, *Agrobacterium*-mediated plant transformation: The biology behind the "gene-jockeying" tool, *Microbiol Mol Biol Rev*, 67 (2003) 16-37.

- 103 Hiei Y, Komari T & Kubo T, Transformation of rice mediated by *Agrobacterium tumefaciens*, *Plant Mol Biol*, 35 (1997) 205-218.
- 104 Abdin M Z, Rehman R U, Israr M, Srivastava P S & Bansal K C, Development of transgenic chicory (*Cichorium intybus* L.), in *In vitro applications in crop improvement*, edited by A Mujeeb *et al* (Science Publisher, Inc., USA) 2004, 285-296.
- 105 Nisha K K, Seetha K, Rajmohan K & Purushothama M G, *Agrobacterium tumefaciens*-mediated transformation of Brahmi [*Bacopa monniera* (L.) Wettst.], a popular medicinal herb of India, *Curr Sci*, 85 1 (2003) 85-89.
- 106 Han K H, Fleming P, Walker K, Loper M, Chilton W S *et al*, Genetic transformation of mature *Taxus*: An approach to genetically control the *in vitro* production of the anticancer drug, taxol, *Plant Sci*, 95 (1994) 187-196.
- 107 Wang H M & To K Y, *Agrobacterium*-mediated transformation in the high-value medicinal plant, *Echinacea purpurea*, *Plant Sci*, 66 (2004) 1087-1096.
- 108 Park S U, Chae Y A & Facchini P J, Genetic transformation of the figwort, *Scrophularia buergeriana* Miq., an Oriental medicinal plant, *Plant Cell Rep*, 21 (2003) 1194-1198.
- 109 Sales E, Segura J & Arrillaga I, *Agrobacterium tumefaciens*-mediated genetic transformation of the cardenolide-producing plant *Digitalis minor* L., *Planta Med*, 69 (2003) 143-147.
- 110 Samanani N, Park S U & Facchini P J, *In vitro* regeneration and genetic transformation of the berberine-producing plant, *Thalictrum flavum* ssp. *glaucum*, *Physiol Plant*, 116 (2002) 79-86.
- 111 Chen D H, Ye H C & Li G F, Expression of a chimeric farnesyl diphosphate synthase gene in *Artemisia annua* L. transgenic plants via *Agrobacterium tumefaciens*-mediated transformation, *Plant Sci*, 155 (2000) 179-185.
- 112 Dai S, Zheng P, Marmey P, Zhang S, Tian W *et al*, Comparative analysis of transgenic rice plants obtained by *Agrobacterium* mediated transformation and particle bombardment, *Mol Breed*, 7 (2001) 25-33.
- 113 Hu T, Metz S, Chay C, Zhou H P, Biest N *et al*, *Agrobacterium* mediated large-scale transformation of wheat (*Triticum aestivum* L.) using glyphosate selection, *Plant Cell Rep*, 21 (2003) 1010-1019.
- 114 Broothaerts W, Mitchel H J, Weir B, Kaines S, Smith L M A *et al*, Gene transfer to plants by diverse species of bacteria, *Nature (Lond)*, 433 (2005) 629-633.
- 115 Hosokawa K, Matsuki R, Oikawa Y & Yamamura S, Production of transgenic gentian plants by particle bombardment of suspension-culture cells, *Plant Cell Rep*, 19 (2000) 454-458.
- 116 Leo A H Z, Christou P & Leech M J, Transformation of the tropane alkaloid-producing medicinal plant *Hyoscyamus muticus* by particle bombardment, *Transgenic Res*, 9 (2000) 163-168.
- 117 Sawahel W, A stable genetic transformation of garlic plants using particle bombardment, *Cell Mol Biol Lett*, 7 (2002) 49-59.
- 118 Chand P K, Ochatt S J, Rech E L, Power J B & Davey M R, Electroporation stimulates plant regeneration from protoplasts of the woody medicinal species, *Solarium dulcamara* L., *J Exp Bot*, 39 (1988) 1267-1274.
- 119 Boynton J E, Gillham N W, Harris E H, Hosler J P, Johnson A M *et al*, Chloroplast transformation in *Chlamydomonas* with high velocity microprojectiles, *Science*, 240 (1988) 1534-1538.
- 120 Daneil H, Kumar S & Dufourmantel N, Breakthrough in chloroplast genetic engineering of agronomically important crops, *Trends Biotechnol*, 23 (2005) 238-245.
- 121 Bansal K C & Sharma R K, Chloroplast transformation as a tool for prevention of gene flow from GM crops to weedy or wild relatives, *Curr Sci*, 84 (2003) 1286.
- 122 Ferreira J F S & Duke S O, Approaches for maximizing biosynthesis of medicinal plant secondary metabolites, *AgBiotech News Inform*, 9 (1997) 309-315.
- 123 Charlwood B V & Pletsch M, Manipulation of natural product accumulation in plants through genetic engineering, *J Herbs Spices Med Plants*, 9 (2002) 139-151.
- 124 Zhang L, Ding R, Chai Y, Bonfill M, Moyano E *et al*, Engineering tropane biosynthetic pathway in *Hyoscyamus niger* hairy root cultures, *Proc Natl Acad Sci USA*, 101 (2004) 6786-6791.
- 125 Yun D, Hashimoto T & Yamada Y, Metabolic engineering of medicinal plants: transgenic *Atropa belladonna* with an improved alkaloid composition, *Proc Natl Acad Sci USA*, 89 (1992) 11799-11803.
- 126 Facchini P J, Park S U, Bird D A & Samanani N, Toward the metabolic engineering of benzyloisoquinoline alkaloid biosynthesis in opium poppy and related species, *Rec Res Dev Phytochem*, 4 (2001) 31-47.
- 127 Chen D H, Ye H C & Li G F, Expression of a chimeric farnesyl diphosphate synthase gene in *Artemisia annua* L. transgenic plants via *Agrobacterium tumefaciens*-mediated transformation, *Plant Sci*, 155 (2000) 179-185.
- 128 Chen D H, Liu C J, Ye H C, Li G F, Liu B Y *et al*, Ri-mediated transformation of *Artemisia annua* with a recombinant farnesyl diphosphate synthase gene for artemisinin production, *Plant Cell Tissue Organ Cult*, 57 (1999) 157-162.
- 129 Dubey T & Guerra D J, Use of biotechnology for growing medicinal plants, *Recent Prog Med Plants*, 5 (2002) 47-61.
- 130 Saito K, Yamazaki M, Anzai H, Yoneyama K & Murakoshi I, Transgenic herbicide-resistant *Atropa belladonna* using an Ri binary vector and inheritance of the transgenic trait, *Plant Cell Rep*, 11 (1992) 219-224.
- 131 Choi Y E, Jeong J H, In J K & Yang D C, Production of herbicide-resistant transgenic *Panax ginseng* through the introduction of the phosphinothricin acetyltransferase gene and successful soil transfer, *Plant Cell Rep*, 21 (2003) 563-568.
- 132 De Donato M, Mozzetti C & Perucco E, Evaluation of atrazine resistance in protoplast regenerants of *Solanum nigrum* L., *Acta Hort*, 280 (1990) 263-269.
- 133 Chen W P & Punja Z K, *Agrobacterium*-mediated transformation of American ginseng with a rice chitinase gene, *Plant Cell Rep*, 20 (2002) 1039-1045.
- 134 Punja Z K & Chen W P, Tissue culture of American ginseng and genetic engineering to express anti fungal proteins, *Acta Hort*, 625 (2003) 395-401.
- 135 Zschocke S, Klaiber I, Bauer R, & Vogler B, HPLC-coupled spectroscopic techniques (UV, MS, NMR) for the structure elucidation of phthalides in *Ligusticum chuanxiong*, *Mol Diversity*, 9 (2005) 33-39.
- 136 Wolfender J, Ndjoko K & Hostettmann K, The potential of LC-NMR in phytochemical analysis, *Phytochem Anal*, 12 (2001) 2-22.
- 137 Schena M, Sharon D, Davis R W & Brown P O, Quantitative monitoring of gene expression patterns with a complementary DNA microarray, *Science*, 270 (1995) 467-470.

- 138 Schena M, Shalon D, Heller R, Chai A, Brown P O & Davis R W, Parallel human genome analysis: Microarray-based expression monitoring of 1000 genes, *Proc Natl Acad Sci USA*, 93 (1996) 10614-10619.
- 139 Taniguchi M, Miura K, Iwao H & Yamanaka S, Quantitative assessments of DNA microarrays comparison with northern blot analyses, *Genomics*, 71 (2001) 34-39.
- 140 Yamamoto M, Wakatsuki T, Hada A & Ryo A, Use of serial analysis of gene expression (SAGE) technology, *J Immunol Methods*, 250 (2001) 45-66.
- 141 Bertelsen A H & Velculescu V E, High-throughput gene expression analysis using SAGE, *Drug Discov Today*, 3 (1998) 152-159.
- 142 Lee Y, Sultana R, Pertea G, Cho J, Karamycheva S *et al*, Cross referencing eukaryotic genomes: TIGR orthologous gene alignments (TOGA), *Genome Res*, 12 (2002) 493-502.
- 143 Sogayar M C, Camargo A A, Bettoni F, Carraro D M, Pires L C *et al*, Ludwig-FAPESP transcript finishing initiative: A transcript finishing initiative for closing gaps in the human transcriptome, *Genome Res*, 14 (2004) 1413-1423.
- 144 Coldren C D, Hashim P, Ali J M, Oh S K, Sinskey A J & Rha C, Gene expression changes in the human fibroblast induced by *Centella asiatica* triterpenoids, *Planta Med*, 69 (2003) 725-732.
- 145 Iizuka N, Oka M, Yamamoto K, Tangoku A, Miyamoto K *et al*, Identification of common or distinct genes related to antitumor activities of a medicinal herb and its major component by oligonucleotide microarray, *Int J Cancer*, 107 (2003) 666-672.
- 146 Christen Y, Olano-Martin E & Packer L, Egb 761 in the postgenomic era: New tools from molecular biology for the study of complex products such as *Ginkgo biloba* extract, *Cell Mol Biol*, 48 (2002) 593-599.
- 147 Watanabe C M, Wolffram S, Ader P, Rimbach G, Packer L & Maguire *et al*, The *in vivo* neuromodulatory effects of the herbal medicine *Ginkgo biloba*, *Proc Natl Acad Sci USA*, 98 (2001) 6577-6580.
- 148 De Feudis F V, Papadopoulos V & Drieu K, *Ginkgo biloba* extracts and cancer: A research area in its infancy, *Fundam Clin Pharmacol*, 17 (2003) 405-417.
- 149 Fugh-Berman A, Herb-drug interactions, *Lancet*, 355 (2000) 355134-138.
- 150 Coxeter P D, McLachlan A J, Duke C C & Roufogalis B D, Herb-drug interactions: An evidence based approach, *Curr Med Chem*, 11 (2004) 1513-1525.
- 151 Joshi K, Chavan P, Warude D & Patwardhan B, Molecular markers in herbal drug technology, *Curr Sci*, 87 (2004) 159-165.
- 152 Warude D, Chavan P, Joshi K & Patwardhan B, Isolation of DNA from fresh and dry samples having highly acidic tissues, *Plant Mol Biol Rep*, 21 (2003) 1-6.
- 153 Carles M, Lee T, Moganti S, Lenigk R, Tsim K W, Ip N Y *et al*, Chips and Qi: microcomponent-based analysis in traditional Chinese medicine, *Fresenius J Anal Chem*, 371 (2002) 190-194.
- 154 Trau D, Lee T M, Lao A I, Lenigk R, Hsing I M *et al*, Genotyping on a complementary metal oxide semiconductor silicon polymerase chain reaction chip with integrated DNA microarray, *Anal Chem*, 74 (2002) 3168-3173.
- 155 Blower P E, Yang C, Fligner M A, Verducci J S, Yu L *et al*, Pharmacogenomic analysis: Correlating molecular substructure classes with microarray gene expression data, *Pharmacogenomics J*, 2 (2002) 259-271.
- 156 Tsoi P Y, Wu H S, Wong M S, Chen S L, Fong W F *et al*, Genotyping and species identification of *Fritillaria* by DNA chip technology, *Acta Pharm Sin*, 4 (2003) 185-190.
- 157 Zhang Y B, Wang J, Wang Z T, But P P & Shaw P C, DNA microarray for identification of the herb of *Dendrobium* species from Chinese medicinal formulations, *Planta Med*, 69 (2003) 1172-1174.
- 158 Kung P C, Peck K, Lee Y, Sher Y & Cheng Y, Phytomics: A genomic-based approach to herbal compositions, *European Pat EP1263988* (to University Yale, USA). 12 November, 2002.
- 159 Breyne P, Dreesen, R, Cannoot B, Rombaut D, Vandepoele K *et al*, Quantitative cDNA-AFLP analysis for genome-wide expression studies, *Mol Genet Genomics*, 269 (2003) 173-179.
- 160 Breyne P & Zabeau M, Genome-wide expression analysis of plant cell cycle modulated genes, *Curr Opin Plant Biol*, 4 (2001) 136-142.
- 161 Yu Z H, Mackill D J, Bonman J M & Tanksley S D, Tagging genes for blast resistance in rice via linkage to RFLP markers, *Theor Appl. Genet*, 81 (1991) 471-476.
- 162 Ma Z Q, Sorrells M E & Tanksley S D, RFLP markers linked to powdery mildew resistance genes *Pm1*, *Pm2*, *Pm3* and *Pm4* in wheat, *Genome*, 37 (1994) 871-875.
- 163 Khanuja, S P S, Paul S, Shasany A K, Gupta A K, Darokar M P *et al*, High artemisinin yielding plant genotype 'CIM-Arogya', *U S Patent 20070089211* (Council of Scientific and Industrial Research, India), 19 April 2007.
- 164 Murata J, Roepke J, Gordon H & Luca V D, The leaf epidermome of *Catharanthus roseus* reveals its biochemical specialization, *Plant Cell*, 20 (2008) 524-542.
- 165 Rischer H, Oresi M, Seppanen-Laakso T, Katajamaa M, Lammertyn F *et al*, Gene-to metabolite networks for terpenoid indole alkaloid biosynthesis in *Catharanthus roseus* cells, *Proc Natl Acad Sci USA*, 103 (2006) 5614-5619.
- 166 Goossens A, Hakkinen S T & Laakso I *et al*, A functional genomics approach toward the understanding of secondary metabolism in plant cells, *Proc Natl Acad Sci USA*, 100 14 (2003) 8595-8600.
- 167 Zulak K G, Cornish A, Daskalchuk T E, Deyholos M K, Goodenowe D B *et al*, Gene transcript and metabolite profiling of elicitor-induced opium poppy cell cultures reveals the coordinate regulation of primary and secondary metabolism, *Planta*, 225 (2005) 1085-1106.
- 168 Berteaa C M, Vostera A, Francel W A, Maffeib V M, Beekwildera J *et al*, Isoprenoid biosynthesis in *Artemisia annua*: Cloning and heterologous expression of a germacrene A synthase from a glandular trichome cDNA library, *Arch Biochem Biophys*, 448 (2006) 3-12.