

All together now, a magical mystery tour of the maize shoot meristem

Qingyu Wu¹, Fang Xu¹ and David Jackson

Crop yield improvement requires optimization of shoot architecture, and can be facilitated by understanding shoot apical meristem (SAM) development. Maize, as one of the most important cereal crops worldwide, is also a model system and has significantly contributed to our fundamental understanding of SAM development. In this review, we focus on recent progress and will discuss communication between different meristem regulators, including CLAVATA receptors and ligands, transcription factors, small RNAs and hormones, as well as the importance of communication between different SAM regions.

Address

Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, United States

Corresponding author: Jackson, David (jacksond@cshl.edu)

¹The first two authors contributed equally to the work.

Current Opinion in Plant Biology 2018, 45:26–35

This review comes from a themed issue on **Cell signalling and gene regulation**

Edited by **Jorge Casal** and **Javier Palatnik**

<https://doi.org/10.1016/j.pbi.2018.04.010>

1369-5266/© 2018 Elsevier Ltd. All rights reserved.

Introduction

The maize shoot system is derived from the shoot apical meristem (SAM), a pool of pluripotent stem cells which have the ability of self-renewal, and initiate new leaves and axillary meristems [1]. Due to its importance in determining shoot architecture and its self-organizing capability, the SAM has brought much attention to scientists for more than a century. As in other plants, the maize SAM is formed during embryogenesis, starting from ~7 days post pollination [2]. In contrast to animals, where organogenesis is usually complete during embryonic development, most new tissues and organs are formed during post-embryonic development in plants [3]. During maize embryogenesis, around five leaves are made before the seed matures and the embryo becomes dormant [4]. Upon germination, the SAM becomes active once more, and new leaves initiate continually until the transition stage, when the vegetative SAM transits to an

inflorescence meristem (IM) that produces spikelets and flowers. Axillary meristems produced in association with leaf primordia can produce branches (tillers), or ear inflorescence shoots.

The SAM is comprised of different zones, based on their cellular activity. At the tip, the stem cell niche contains a pool of slowly dividing pluripotent stem cells, and is named the central zone (CZ). Surrounding the CZ is the peripheral zone (PZ), where cells divide more rapidly, and will generate leaf or axillary meristem primordia. Lastly, the rib zone (RZ) lies below the CZ, and cells dividing there will form the stem. A small group of cells called the organizing center (OC), sitting between the CZ and RZ was defined more recently based on molecular markers, and is important for communication between different zones (reviewed in [5,6]). A pioneering study showed that in some plants (though not in maize), the SAM can regenerate after the vast majority of cells are surgically removed [7]. This suggests that the SAM is not rigidly programmed, but can reestablish different zones and communication pathways to rebuild the structure of the meristem. Stem cell activity is at the heart of the SAM, and when this is disrupted by specific mutations, plant growth will be abnormal or will cease completely. In contrast, mutants that upregulate stem cell activity lead to bigger meristems with abnormally flattened and split stems and inflorescences, called fasciation. Such mutants have significantly contributed to our understanding of signaling and communication between zones of the maize SAM (summarized in Table 1). Recent progress has been rapid, fueled by advances in genomics, however the picture is still far from complete. In this review, we will synthesize findings from recent studies, focusing on receptor based communication between different zones of the maize SAM and intracellular signal transduction.

The classical CLAVATA (CLV)-WUSCHEL (WUS) model established in *Arabidopsis* is largely conserved in maize



The CLV-WUS model has long been recognized as a key feedback pathway that regulates communication between different zones in the *Arabidopsis* SAM [1]. It relies on communication between a series of receptors, peptide ligands and transcription factors that are expressed in different zones. Central to this complex network is WUS, a homeodomain transcription factor expressed in the OC to promote stem cell fate [8], and CLV3, a small peptide ligand that is secreted from cells in

Table 1

Summary of fasciated inflorescence mutants. Scanning electron microscopy images of wild-type and fasciated mutant ear primordia are presented, except for *Cg1*, for which a tassel primordium is shown. Inflorescence meristems are shaded in yellow. Protein structures are illustrated with indicated domains except for *Cg1*, a miRNA, for which transcript is illustrated. The references for the mutants are listed under their gene IDs. SP: signal peptide; LRR: leucine-rich repeat; TM: transmembrane; KD: kinase domain; bZIP: basic leucine zipper; CLE: Clavata3/Embryo Surrounding Region (ESR)-Related; SBP: SQUAMOSA promoter Binding Protein; SNH: Synovial Translocation (SYT) N-terminal Homology; PAZ: Piwi Argonaut and Zwiile; dsRB: double strand RNA Binding

| | IM | SAM size | Protein structure | <i>Arabidopsis</i> ortholog |
|--|----|----------------|-------------------|--|
| WT | | N/A | N/A | N/A |
| <i>fea2</i> GRMZM2G104925 (14) | | Enlarged | | CLV2 (AT1g65380) |
| <i>fea3</i> GRAZM2G166524 (19) | | Enlarged | | AtFEA3 (AT3g25670) |
| <i>fea4</i> GRMZM2G133331 (47) | | Enlarged | | PAN (AT1g68640) |
| <i>td1</i> GRMZM2G300133 (15) | | Not determined | | CLV1 (AT1g75820) |
| <i>ct2</i> GRMZM2G064732 (21) | | Enlarged | | GPA1 (AT2G26300) |
| <i>Zmfc1</i> GRMZM2G165836 (19) | | Enlarged | | CLE27 (AT3G25905) |
| <i>ub2,ub3</i> GRMZM2G160917 GRMZM2G460544 (42) | | Not determined | | AtSPL9 (AT2G42200) AtSPL15 (AT3G57920) |
| <i>gif1</i> GRMZM2G180246 (45) | | Not determined | | AtGIF1 (AT5G28640) |
| <i>fzt1</i> GRMZM2G040762 (65) | | Not determined | | DCL1 (AT1G01040) |
| <i>Cg1</i> GRMZM2G022489 (67) | | Not determined | | miR156B (AT4G30972) miR156C (AT4G31877) |

Table 1 (Continued)

| | IM | SAM size | Protein structure | <i>Arabidopsis</i> ortholog |
|---------------------------------------|---|----------|--|-----------------------------|
| <i>Zmcrn</i> GRMZM2G032132 (28) |  | Enlarged |  | CRN (AT5G13290) |

the CZ and perceived by leucine-rich repeat receptor-like kinases (LRR-RLKs), such as CLV1, and a leucine-rich repeat receptor-like protein (LRR-RLP) CLV2, resulting in the repression of WUS transcription [9–11]. Key to this feedback loop is the precise spatial expression of these components and the movement of CLV3 and WUS. CLV3 is secreted, and presumably moves through the extracellular matrix by diffusion, though the peptide has never been localized *in vivo*, and WUS protein moves cell to cell through plasmodesmata [12,13]. The CLV3 receptor *CLV1* is expressed below the CZ and surrounding the WUS domain, while *CLV2* is expressed broadly in the SAM and throughout the plant.

Conservation of the CLV-WUS pathway in maize is evident through studies using ‘*fasciated ear*’ (*fea*) and similar mutants (Figure 1). For example, *FASCATED EAR2* (*FEA2*) encodes an LRR-RLP that is orthologous to *CLV2*, and *fea2* mutants have enlarged vegetative SAMs and IMs [14]. Further evidence of conservation came from the cloning of *THICK TASSEL DWARF1* (*TD1*), which encodes a *CLV1* ortholog [15]. *td1* mutants have over-proliferating IMs, similar to *fea2* [15]. While additional LRR-RLKs, such as BARELY ANY MERISTEMs (BAMs), function redundantly with *CLV1* in *Arabidopsis* [16,17], it remains unclear whether the same is true for maize. Notably, *TD1* expression is similar to *BAM* genes, with expression high in the peripheral region of the vegetative SAM and in leaf primordia, and absent from the CZ [16,17]. More work is needed to understand the intricacies of the TD1-BAM clade in maize meristem regulation. Unlike the clearly defined *CLV1* and *CLV2* orthologs, maize has two *WUS* orthologs, *ZmWUS1* and *ZmWUS2*, however both are expressed at very low levels in the vegetative SAM [18]. However, *ZmWUS1* is clearly expressed in the OC of the inflorescence SAM, similar to *Arabidopsis* [19**]. *Zea mays* CLAVATA3/EMBRYO SURROUNDING REGION-RELATED7 (*ZmCLE7*) and *ZmCLE14* are candidates for maize *CLV3* orthologs, based on phylogenetic analysis [19**,20*], and treatment with these peptides affects SAM size in a predictable manner [19**], but functional characterization of *WUS* and *CLV3* orthologs is necessary to understand their true role in maize.


Signaling immediately downstream of the CLV receptors is poorly understood, possibly due to genetic redundancy or pleiotropy caused by converging signaling

pathways. However, analysis of maize *compact plant2* (*ct2*) mutants, which also have fasciated ears and enlarged SAMs, reveals a role of heterotrimeric G-proteins in meristem regulation [21]. *ct2* mutants are partially resistant to CLV3 peptide treatment and *CT2*, which encodes $G\alpha$, is in a protein complex with *FEA2*, suggesting that *CT2* and *FEA2* together transmit the CLV signal. Heterotrimeric G-protein signaling also appears to be important in *Arabidopsis* meristem regulation, where the heterotrimeric G protein beta subunit, *AGB1* interacts with another CLV-like receptor RECEPTOR-LIKE PROTEIN KINASE2 (RPK2) to control SAM size [22], but in the case of $G\alpha$ is more redundant, since an increase in SAM size is only seen in quadruple mutants of $G\alpha$ with 3 related ‘*eXtra-Large G protein*’ genes [23]. Another downstream signaling component of CLV receptors is CORYNE (CRN), a transmembrane receptor-like pseudo-kinase [24–26]. In *Arabidopsis*, CRN directly associates with *CLV2*, though how this interaction mediates signal transduction remains unclear, since CRN lacks kinase activity [26,27]. A recent study shows that its maize ortholog, *ZmCRN*, also functions downstream of *FEA2*, the *CLV2* ortholog. This study shows that *FEA2* transmits signals from two distinct CLE peptides, *ZmCLE7* and *ZmFCP1* through specific interactions with *CT2* and *ZmCRN*, respectively [28*]. This finding brings new insights into how LRR-RLPs can transmit different peptide signals through associating with different downstream signaling components. However details of these pathways are lacking, for example *FEA2* is an LRR-RLP with no intracellular signaling domain, and *fea2* mutants are insensitive to treatments using several different CLE peptides [19**], so it is likely that *FEA2* works as a co-receptor of multiple LRR-RLK receptors. The co-receptor idea is also based on results with *CLV2*, which cannot bind CLE peptides directly [29], so it will be interesting to identify the co-receptors that associate with *FEA2* and bind CLE peptides (Figure 1).

Signaling from stem cell descendants back to the stem cell niche

In addition to the CLV-WUS pathway, which enables communication between cells within the SAM, recent studies in maize further highlight that stem cell descendants and initiating leaf primordia also send signals that feedback on SAM regulation. For example, the maize *YABBY* genes *DROOPING LEAF1* and *DROOPING*

Figure 1



Signal transduction in maize shoot meristems. In the center of the SAM, a putative CLV3 ortholog, ZmCLE7, interacts with FEA2 and a putative CLV1 ortholog, and the heterotrimeric G protein α subunit CT2 mediates CLV signaling by coordinating with FEA2. KN1 also functions in SAM maintenance, acting redundantly with RS1, and forms heterodimers with BLH12 or BLH14. Meanwhile, on the SAM periphery, FEA3 and a FEA2-CRN complex perceive signals from ZmFCP1, while the ligand of TD1 remains unknown. Signaling via CLV receptors leads to the repression of the stem cell promoter *ZmWUS*. Also on the SAM periphery, FEA4 interacts with MSCA1, and GIF1 regulates the expression of *UB3*, to inhibit stem cell maintenance. Lastly, acting from the leaf primordia, YABBY transcription factors promote SAM maintenance by an unknown signal.

LEAF2 (*DRL1, 2*) were initially characterized due to their redundant effects on leaf angle, an important agronomic trait. *DRL1* and *2* are expressed exclusively in leaf primordia, yet the double mutants have smaller SAMs, and similar findings in *Arabidopsis* also suggest that signals from organ primordia non-autonomously regulate SAM development [30,31], however, the signaling mechanism remains elusive. Another example comes from the characterization of *fasciated ear3* (*fea3*) mutants. *FEA3* encodes an LRR-RLP similar to FEA2, but unlike other *CLV* genes that are expressed in stem cells and above and around the OC, *FEA3* is expressed within and below the OC, and in leaf primordia. *FEA3* is involved in perception of a signal from the CLE peptide ZmFCP1, expressed in PZ and leaf primordia, and *FEA3* inhibits the expansion of *ZmWUS1* downwards below the OC. Consistently, both *fea3* and *Zmfcf1* mutants have bigger SAMs and IMs, and *fea3* is epistatic to *Zmfcf1* with respect to SAM size [19]. Therefore, *FEA3* and ZmFCP1 define a new CLV feedback that appears to

be conserved in *Arabidopsis* [19], but a major question remains in the CLE field whether the CLE peptides move, as predicted, and whether it is by passive diffusion or active transport?

Additional transcription factors regulate the maize SAM

Besides *WUS*, additional transcription factors expressed in different SAM zones play important roles in maize development. One of the best studied is *KNOTTED1* (KN1), the founding member of the KN1-related homeobox (KNOX) family [32–35]. Severe KN1 loss-of-function mutants fail to initiate or maintain the SAM, although in some genetic backgrounds somewhat normal plants survive albeit with smaller inflorescence and vegetative SAMs [32,34]. *KN1* is expressed throughout the SAM but excluded from leaf initiation sites, leaf primordia and the L1 layer [36], although KN1 protein traffics cell to cell through plasmodesmata into the L1, providing a possible mechanism to coordinate development of different cell

layers [37,38]. A recent study shows that KN1 forms heterodimers with BEL1-like homeobox transcription factors BLH12 and BLH14 [39]. *blh12,blh14* double mutants make slightly shorter but otherwise normal SAMs, but axillary meristems were not maintained, similar to the case in *kn1* mutants [39]. With the expanding use of transcriptome and ChIP-seq profiling, it was found that KN1 directly targets many developmental regulators, including transcription factors and genes participating in auxin and gibberellic acid hormonal pathways, revealing a complicated regulatory hierarchy for SAM development controlled by KN1 [40*,41]. However, the interaction between KN1 and ZmWUS still remains obscure, and seeking the common targets of both proteins could help to gain a better understanding of this question.

A major limitation of genetic analysis in plants is redundancy, as highlighted in a recent report [40*]. Expression of the *KNOX* gene *ROUGH SHEATH1 (RS1)* overlaps that of *KN1* [40*], but *rs1* mutants have no obvious phenotype. However in double mutants, *rs1* exacerbates several aspects of the *kn1* phenotype. ChIP-seq experiments show that RS1 shares many *in vivo* binding targets with KN1, suggesting unequal redundancy between *KNOX* family members occurs at the level of transcriptional outputs. Overcoming genetic redundancy was also necessary to demonstrate that inflorescence development is regulated by SQUAMOSA PROMOTER BINDING (SBP)-box transcription factors UNBRANCHED2 (UB2) and UB3 [42**]. *ub2,3* double mutants have enlarged IMs and fasciated ears, and both genes are expressed throughout the periphery of the meristem. Previous studies raised the idea that weak alleles of fasciated mutants could enhance kernel row number (KRN) [19**,43**], an important agronomic trait, and fine-mapping of a major KRN quantitative trait locus (QTL), *KRN4*, found that a transposon insertion ~ 60 kb downstream of the *UB3* gene regulates its expression, and is responsible for maize KRN variation [44**]. More recently, *UB3* was found to be a transcriptional target of GROWTH-REGULATING FACTOR (GRF)-INTERACTING FACTOR 1 (GIF1) (Figure 1), and its expression is reduced in *gif1* mutants [45*]. Similar to *ub2,3* mutants, tassels and ears are fasciated in the *gif1* mutants. *UB3* itself may function by regulating cytokinins, as it binds the promoters of *LONELY GUY1 (LOG1)* and *TYPE-A RESPONSE REGULATOR (ARR)* genes that are involved in cytokinin biosynthesis and signaling [46].

Another gene critical to maize SAM regulation is *FASCIATED EAR4 (FEA4)*, which encodes a bZIP transcription factor orthologous to *Arabidopsis PERIANTHIA (PAN)* [47*]. *fea4* mutants have enlarged vegetative SAMs and very severely fasciated IMs, and double mutant analyses indicate that it functions as a negative regulator of meristem size in parallel to CLV-WUS signaling. However, *PAN* was previously reported as a controller of floral organ

patterning but not meristem size [48], revealing the importance of studying basic processes in different model systems in order to reveal gene functions. Like *UB2* and *UB3*, *FEA4* is expressed in the SAM periphery, again highlighting the possibility that signals feedback from the PZ to the stem cell niche (Figure 1). *FEA4* may be regulated by redox signaling in the SAM, since it interacts with a glutaredoxin protein MALE STERILE CONVERTED ANTHER1 (*MSCA1*) [49]. *mzca1* loss-of-function mutants have smaller SAMs, whereas *MSCA1* misexpression in dominant *MSCA1-Abserrant phyllotaxy2* mutants leads to a bigger SAM and decussate phyllotaxy. The importance of redox signaling in the SAM was also recently discovered in *Arabidopsis*, where redox treatments and mutants alter SAM size [50]. Although it is confusing that so many factors control SAM size, genomic analysis can help clarify their relationships. In one example, ChIP-seq combined with RNA-seq profiling revealed that *FEA4* and *KN1* oppositely regulate a common set of target genes, suggesting that *FEA4* functions in part in opposition to *KN1* [47*]. A remaining question is how does *MSCA1* interaction and redox signaling affect the function of *FEA4*?

Hormones crosstalk to regulate SAM development

Plant hormone signaling regulates development and communication between different regions of the SAM, converging with the pathways discussed earlier. The most well studied hormones are cytokinin and auxin, with cytokinin promoting meristem proliferation and auxin driving lateral primordium initiation. Insights have come from characterization of maize mutants with altered auxin biosynthesis, for example, the *sparse inflorescence1 (spi1)* mutants that are defective in auxin biosynthesis have *pin*-like inflorescence meristems [51]. *SPI1* encodes a flavin monooxygenase similar to the *YUCCA* genes of *Arabidopsis*, which are involved in local auxin biosynthesis. Another example is *vanishing tassel2 (vt2)*. *VT2* encodes an ortholog of tryptophan aminotransferase of *Arabidopsis (TAA1)* that is also required for auxin biosynthesis [52]. *spi1* and *vt2* single mutants have reduced auxin levels, and the tassels are smaller and barren, lacking lateral branches and functional spikelets. The reduction in auxin levels in *spi1,vt2* double mutants is similar to single mutants, suggesting they work in a common pathway. Importantly, the finding that *VT2* and *SPI1* are epistatic led to a new hypothesis that *TAA1* and *YUCCA* function in the same IAA biosynthetic pathway, which was later confirmed in *Arabidopsis* [53].

Analysis of additional *barren* mutants revealed important roles for auxin signaling and transport. *BARREN INFLORESCENCE1 (BIF1)* and *BIF4* encode AUXIN/INDOLE-3-ACETIC ACID (Aux/IAA) proteins, key players in auxin signaling [54]. *BIF1* and *BIF4* control inflorescence axillary meristem initiation by regulating

the expression of *BARREN STALK1 (BA1)*, a basic helix–loop–helix transcription factor [55]. In addition, *BARREN INFLORESCENCE2 (BIF2)* encodes a co-ortholog of the serine/threonine protein kinase PINOID (PID), and regulates auxin transport through affecting the subcellular localization of a PINFORMED1 auxin efflux transporter ortholog (ZmPIN1a) during maize inflorescence development [56,57]. The importance of auxin transport was also revealed by studies of the auxin influx carrier mutant, *Zmaux1*, which reduces tassel branch number and has fewer spikelets [58]. A study of the role of natural variation in meristem size control further showed that expression of its paralog, *ZmLAX2*, associates with SAM morphological diversity in a genome-wide association study (GWAS) [59^{*}]. *In situ* hybridization revealed that *ZmLAX2* transcript accumulation coincided with PIN1 localization, suggesting that the balance between auxin efflux and influx needs to be balanced for proper SAM development, and that changes in *ZmLAX2* expression may be responsible for changes in SAM size [59^{*}]. Earlier work also identified *ABERRANT PHYLLLOTAXY1 (ABPH1)* as a cytokinin-inducible type A response regulator, indicating that cytokinin controls SAM size and phyllotaxy [60,61]. *ABPH1* expression is dependent on auxin transport, and it activates *PIN1* expression [61] and therefore orchestrates phyllotaxy by restricting SAM size through negative regulation of a cytokinin signal and promoting leaf initiation by positively regulating auxin signaling, revealing a complex interaction between auxin and cytokinin in the SAM [61].

Additional pathways required for maize shoot meristem maintenance

Certain metabolites and mineral nutrients are also important for SAM maintenance. Mutations in a thiamine biosynthetic gene, *THIAMINE BIOSYNTHESIS2 (THI2)*, result in defects in meristem maintenance and a progressive reduction in SAM size that leads to premature shoot abortion, providing a new perspective on metabolic requirements of the SAM [62]. Mutations in *TASSEL-LESS1 (TLS1)*, which encodes an aquaporin family protein that facilitates the transport of boron also lead to similar defects in vegetative and inflorescence development [63]. Boron content is reduced in *tls1* mutants and application of boron can rescue the mutant phenotype, highlighting the importance of this mineral in meristem function. The recently characterized *narrow odd dwarf (nod)* mutants provide new insights into how cell expansion and division affects meristem development [64]. *NOD* encodes CELL NUMBER REGULATOR13 (CNR13) [64], and *nod* mutants have pleiotropic defects, including fewer and smaller leaf cells, dwarf and bushy shoots, reduced SAM size, barren tassels, as well as small, partially fertile ears. Remarkably, many of the differentially expressed genes in *nod* mutants overlap with KN1 targets, suggesting crosstalk between these two genes. Small RNAs, key players in many plant

processes have also been implicated in meristem maintenance in maize. For instance, *FUZZY TASSEL (FZT)* encodes a homolog of DICER-LIKE1, a key enzyme in microRNA biogenesis. While null alleles are lethal, weak alleles are defective in the production of a subset of miRNAs, and exhibit fasciated inflorescences [65]. Overexpression of certain miRNAs also resulted in abnormal IM development. For example, the dominant mutant *Corngrass1 (Cg1)*, caused by the overexpression of miR156, has a fasciated inflorescence meristem similar to *fzt*; however, their genetic relationship remains obscure as miRNA156s level are not significantly changed in *fzt* mutants [66]. Remarkably, *Cg1* mutants have more severe phenotypes than plants overexpressing miR156 in *Arabidopsis*, suggesting that the buffering networks of miRNAs are more redundant in *Arabidopsis* [67,68]. Together, these studies suggest that the miRNA accumulation needs to be delicately controlled to orchestrate the expression of their target genes, such as SBP-box transcription factors, which are key regulators in meristem development [66].

Axillary meristems and tillering

Tillers are secondary shoots derived from axillary meristems produced at the base of each leaf [69,70]. Domestication of modern maize from the highly tillering ancestor, teosinte, strongly selected against tillering to enable high planting density and yield [71]. Among the major selected loci were *TEOSINTE BRANCHED1 (TB1)*, which encodes a TB1-CYCLOIDEA-PROLIFERATING CELL FACTOR (TCP) transcription factor that acts as a repressor of axillary bud growth [72], and *GRASSY TILLERS1 (GT1)* [73]. *GT1* encodes an HD-ZIP I protein, and its expression is under the transcriptional control of TB1 and is dramatically reduced in *tb1* mutants. Recent characterization of a third mutant, *tassels replace upper ears1 (tru1)* provides a new molecular mechanism for how TB1 transformed teosinte into modern maize [74^{*}]. *tru1* mutants cause maize to revert back to its highly branched ancestral state, much like *tb1*. *TRU1* encodes an ankyrin repeat protein and is expressed at the base of lateral organs during both vegetative and inflorescence development, but is absent from meristems. Genetic and molecular analyses suggest that TB1 mediates suppression of axillary branches through direct activation of TRU1. Future work is needed to elucidate the downstream signaling pathways for these newly identified domestication targets.

Perspective

With the awesome power of genetics, which will no doubt continue to be the cornerstone for meristem research, a subset of key regulators has been identified in different plant species. Many key signaling pathways including CLV-WUS, KNOX, and plant hormones have conserved functions in maize and dicot species. Yet, the relatively

large meristem size and high genetic diversity make maize a powerful model, and can enable many new discoveries. For instance, the identification of FEA3 and its ligand leads to the discovery of new CLV feedback, which was missed in *Arabidopsis* genetic screens. The functional analysis of FEA4 highlighted its role in meristem regulation, with a much stronger effect than its ortholog in *Arabidopsis*, and led to the discovery of a new pathway acting in parallel with CLV-WUS. Despite the efforts of maize geneticists, many maize fasciation mutants have not yet been characterized, suggesting that meristem signaling pathways are more easily disrupted in maize than in *Arabidopsis*. One possible explanation for this is that maize has been subjected to human selection from ~10 000 years ago [71], while *Arabidopsis* is a wild species. One of the major targets for human selection was for larger ears, which is correlated with bigger meristems [43**]. Thus, the alleles that provide redundancy and buffering of meristem size may have been selected against in modern maize cultivars, providing a great opportunity to use maize as a model system to identify novel meristem regulators.

Genetic redundancy may have hindered discoveries of redundant components in the past, but with the advance of gene editing techniques, it is routine to generate multiplex knockouts. Therefore, progress in addressing many unanswered questions in meristem regulation will speed up in the next decade. For example, the complexity of receptor complexes for recognition of CLE peptides remains to be explored, and *in vivo* localization of CLEs is still missing. So far, only one CLE, ZmFCP1, has been functionally investigated in maize. Additional CLEs and their crosstalk need to be studied. In addition, very little is known about downstream signaling and how the signal is transduced from the membrane to the nucleus. Moreover, it will be interesting to learn more about the interaction between KNOX and CLV-WUS pathways, for example do they share common targets and how do they coordinate? Studies of SAM regulation in maize not only deepen our understanding of basic biological questions, but also enable potential application in agriculture. Our ancestors have successfully selected spontaneous mutations in the key SAM regulators, such as *WUS* and *CLV*, to domesticate wild species into modern cultivars with favorable traits and higher yield [75,76]. Advancing genomic and genome editing technologies may allow us to identify more SAM regulators and precisely fine-tune them to further boost crop yields. Recent work found that CRISPR mutagenesis of tomato *CLV3* cis elements could enhance yield [77**], and engineering of FEA2 and FEA3 by generation of weak alleles increased maize seed production and yield [19**,43**]. All of these examples suggest that the tremendous efforts that have been made toward understanding the SAM development are paying back and could help feed a growing world population.

Acknowledgements

We acknowledge the funding from Agriculture and Food Research Initiative competitive grants 2015-06319 and 2017-06299 of the USDA National Institute of Food and Agriculture, National Science Foundation (NSF) grants 1238202 and 1546837, and HFSP Long-Term fellowship LT000227-2016. We apologize to colleagues whose works were not discussed in this review due to length constraints.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Somssich M, Je BI, Simon R, Jackson D: **CLAVATA-WUSCHEL signaling in the shoot meristem.** *Development* 2016, **143**:3238-3248.
 2. Randolph LF: **Developmental morphology of the caryopsis in maize.** *J Agric Res* 1936, **53**:0881-0916.
 3. Heidstra R, Sabatini S: **Plant and animal stem cells: similar yet different.** *Nat Rev Mol Cell Biol* 2014, **15**:301-312.
 4. Evans MMS, Passas HJ, Poethig RS: **Heterochronic effects of glossy15 mutations on epidermal-cell identity in maize.** *Development* 1994, **120**:1971-1981.
 5. Holt AL, van Haperen JMA, Groot EP, Laux T: **Signaling in shoot and flower meristems of *Arabidopsis thaliana*.** *Curr Opin Plant Biol* 2014, **17**:96-102.
 6. Jackson D: **Vegetative shoot meristems.** In *Handbook of Maize — Its Biology*. Edited by Bennetzen J, Hake S. Springer; 2009:1-12.
 7. Sussex IM: **Developmental programming of the shoot meristem.** *Cell* 1989, **56**:225-229.
 8. Schoof H, Lenhard M, Haecker A, Mayer KFX, Jurgens G, Laux T: **The stem cell population of *Arabidopsis* shoot meristems is maintained by a regulatory loop between the CLAVATA and WUSCHEL genes.** *Cell* 2000, **100**:635-644.
 9. Clark SE, Williams RW, Meyerowitz EM: **The CLAVATA1 gene encodes a putative receptor kinase that controls shoot and floral meristem size in *Arabidopsis*.** *Cell* 1997, **89**:575-585.
 10. Fletcher LC, Brand U, Running MP, Simon R, Meyerowitz EM: **Signaling of cell fate decisions by CLAVATA3 in *Arabidopsis* shoot meristems.** *Science* 1999, **283**:1911-1914.
 11. Jeong S, Trotochaud AE, Clark SE: **The *Arabidopsis* CLAVATA2 gene encodes a receptor-like protein required for the stability of the CLAVATA1 receptor-like kinase.** *Plant Cell* 1999, **11**:1925-1933.
 12. Daum G, Medzihradsky A, Suzuki T, Lohmann JU: **A mechanistic framework for noncell autonomous stem cell induction in *Arabidopsis*.** *Proc Natl Acad Sci U S A* 2014, **111**:14619-14624.
 13. Yadav RK, Perales M, Gruel J, Girke T, Jonsson H, Reddy GV: **WUSCHEL protein movement mediates stem cell homeostasis in the *Arabidopsis* shoot apex.** *Genes Dev* 2011, **25**:2025-2030.
 14. Taguchi-Shiobara F, Yuan Z, Hake S, Jackson D: **The fasciated ear2 gene encodes a leucine-rich repeat receptor-like protein that regulates shoot meristem proliferation in maize.** *Genes Dev* 2001, **15**:2755-2766.
 15. Bommert P, Lunde C, Nardmann J, Vollbrecht E, Running M, Jackson D, Hake S, Werr W: **thick tassel dwarf1 encodes a putative maize ortholog of the *Arabidopsis* CLAVATA1 leucine-rich repeat receptor-like kinase.** *Development* 2005, **132**:1235-1245.
 16. Dievart A, Dalal M, Tax FE, Lacey AD, Huttly A, Li JM, Clark SE: **CLAVATA1 dominant-negative alleles reveal functional overlap between multiple receptor kinases that regulate meristem and organ development.** *Plant Cell* 2003, **15**:1198-1211.

17. Nimchuk ZL, Zhou Y, Tarr PT, Peterson BA, Meyerowitz EM: **Plant stem cell maintenance by transcriptional cross-regulation of related receptor kinases.** *Development* 2015, **142**:1043-1049.
18. Nardmann J, Werr W: **The shoot stem cell niche in angiosperms: expression patterns of WUS orthologues in rice and maize imply major modifications in the course of mono- and dicot evolution.** *Mol Biol Evol* 2006, **23**:2492-2504.
19. Je BI, Gruel J, Lee YK, Bommert P, Arevalo ED, Eveland AL, Wu Q, Goldshmidt A, Meeley R, Bartlett M *et al.*: **Signaling from maize organ primordia via FASCIATED EAR3 regulates stem cell proliferation and yield traits.** *Nat Genet* 2016, **48**.
- The authors identified a new receptor-like protein in stem cell control, FEA3, which perceives the FCP1 peptide signal from differentiating cells. This research reveals a fundamentally new meristem regulation pathway from differentiating cells to the stem cell niche.
20. Goad DM, Zhu CM, Kellogg EA: **Comprehensive identification and clustering of CLV3/ESR-related (CLE) genes in plants finds groups with potentially shared function.** *New Phytol* 2017, **216**:605-616.
- The authors used a new type of analysis to construct a phylogenetic tree for all CLE genes in multiple plants, providing insights for future evolutionary and functional studies.
21. Bommert P, Je BI, Goldshmidt A, Jackson D: **The maize Ga gene COMPACT PLANT2 functions in CLAVATA signalling to control shoot meristem size.** *Nature* 2013, **502**:555-558.
22. Ishida T, Tabata R, Yamada M, Aida M, Mitsumasu K, Fujiwara M, Yamaguchi K, Shigenobu S, Higuchi M, Tsuji H *et al.*: **Heterotrimeric G proteins control stem cell proliferation through CLAVATA signaling in Arabidopsis.** *Embo Rep* 2014, **15**:1202-1209.
23. Urano D, Maruta N, Trusov Y, Stoian R, Wu QY, Liang Y, Jaiswal K, Thung L, Jackson D, Botella JR *et al.*: **Saltational evolution of the heterotrimeric G protein signaling mechanisms in the plant kingdom.** *Sci Signal* 2016, **9**.
24. Miwa H, Betsuyaku S, Iwamoto K, Kinoshita A, Fukuda H, Sawa S: **The receptor-like kinase SOL2 mediates CLE signaling in Arabidopsis.** *Plant Cell Physiol* 2008, **49**:1752-1757.
25. Muller R, Bleckmann A, Simon R: **The receptor kinase CORYNE of Arabidopsis transmits the stem cell-limiting signal CLAVATA3 independently of CLAVATA1.** *Plant Cell* 2008, **20**:934-946.
26. Nimchuk ZL, Tarr PT, Meyerowitz EM: **An evolutionarily conserved pseudokinase mediates stem cell production in plants.** *Plant Cell* 2011, **23**:851-854.
27. Bleckmann A, Weidtkamp-Peters S, Seidel CAM, Simon R: **Stem cell signaling in Arabidopsis requires CRN to localize CLV2 to the plasma membrane.** *Plant Physiol* 2010, **152**:166-176.
28. Je BI, Xu F, Wu Q, Liu L, Meeley R, Gallagher JP, Corcilius L, Payne RJ, Bartlett ME, Jackson D: **The CLAVATA receptor FASCIATED EAR2 responds to distinct CLE peptides by signaling through two downstream effectors.** *Elife* 2018, **7**.
- By identification and characterization of a new fasciated ear mutant, *Zmcrn* in maize, the authors showed that ZmCRN functions as a downstream component for FEA2, further revealing the conservation of the CLV pathway between dicots and monocots. Through genetic analyses, protein interaction tests and peptide assays, the authors showed that FEA2 transmits signaling from two distinct CLE peptides, ZmCLE7 and ZmFCP1, through two different downstream components, CT2 and ZmCRN, respectively. This study provides a novel framework to understand how diverse signaling peptides can activate different downstream pathways through common receptor proteins and bring new insights to understand the signaling specificity by different ligands.
29. Shinohara H, Matsubayashi Y: **Reevaluation of the CLV3-receptor interaction in the shoot apical meristem: dissection of the CLV3 signaling pathway from a direct ligand-binding point of view.** *Plant J* 2015, **82**:328-336.
30. Goldshmidt A, Alvarez JP, Bowman JL, Eshed Y: **Signals derived from YABBY gene activities in organ primordia regulate growth and partitioning of Arabidopsis shoot apical meristems.** *Plant Cell* 2008, **20**:1217-1230.
31. Strable J, Wallace JG, Unger-Wallace E, Briggs S, Bradbury P, Buckler ES, Vollbrecht E: **Maize YABBY genes drooping leaf1 and drooping leaf2 regulate plant architecture.** *Plant Cell* 2017, **29**:1622-1641.
- The authors identified two YABBY transcription factors that regulate leaf angle, a key agronomic trait. Both are expressed in the leaf primordia, and SAM size is reduced in the double mutants.
32. Kerstetter RA, LaudenciaChingcuanco D, Smith LG, Hake S: **Loss-of-function mutations in the maize homeobox gene, knotted1, are defective in shoot meristem maintenance.** *Development* 1997, **124**:3045-3054.
33. Long JA, Moan EI, Medford JI, Barton MK: **A member of the KNOTTED class of homeodomain proteins encoded by the STM gene of Arabidopsis.** *Nature* 1996, **379**:66-69.
34. Vollbrecht E, Reiser L, Hake S: **Shoot meristem size is dependent on inbred background and presence of the maize homeobox gene, knotted1.** *Development* 2000, **127**:3161-3172.
35. Vollbrecht E, Veit B, Sinha N, Hake S: **The developmental gene knotted-1 is a member of a maize homeobox gene family.** *Nature* 1991, **350**:241-243.
36. Jackson D, Veit B, Hake S: **Expression of maize KNOTTED1 related homeobox genes in the shoot apical meristem predicts patterns of morphogenesis in the vegetative shoot.** *Development* 1994, **413**:405-413.
37. Kim JY, Yuan ZA, Cilia M, Khalfan-Jagani Z, Jackson D: **Intercellular trafficking of a KNOTTED1 green fluorescent protein fusion in the leaf and shoot meristem of Arabidopsis.** *Proc Natl Acad Sci U S A* 2002, **99**:4103-4108.
38. Xu XM, Wang J, Xuan ZY, Goldshmidt A, Borrill PGM, Hariharan N, Kim JY, Jackson D: **Chaperonins facilitate KNOTTED1 cell-to-cell trafficking and stem cell function.** *Science* 2011, **333**:1141-1144.
39. Tsuda K, Abraham-Juarez M-J, Maeno A, Dong Z, Aromdee D, Meeley R, Shiroishi T, Nonomura K-i, Hake S: **KNOTTED1 cofactors, BLH12 and BLH14, regulate internode patterning and vein anastomosis in maize.** *Plant Cell* 2017, **29**:1105-1118.
40. Bolduc N, Tyers RG, Freeling M, Hake S: **Unequal redundancy in maize knotted1 homeobox genes.** *Plant Physiol* 2014, **164**:229-238.
- RS1* is a *KNOX* family gene, but *rs1* loss-of-function mutants do not have a phenotype. However, they exacerbate the phenotype of *kn1* mutants, suggesting an unequal redundancy between *KNOX* genes.
41. Bolduc N, Yilmaz A, Mejia-Guerra MK, Morohashi K, O'Connor D, Grotewold E, Hake S: **Unraveling the KNOTTED1 regulatory network in maize meristems.** *Genes Dev* 2012, **26**:1685-1690.
42. Chuck GS, Brown PJ, Meeley R, Hake S: **Maize SBP-box transcription factors unbranched2 and unbranched3 affect yield traits by regulating the rate of lateral primordia initiation.** *Proc Natl Acad Sci U S A* 2014, **111**:18775-18780.
- The authors revealed that the duplicate SBP-box transcription factors UB2 and UB3 function together to regulate the rate of lateral primordia initiation, and the double mutants developed fasciated IMs.
43. Bommert P, Nagasawa NS, Jackson D: **Quantitative variation in maize kernel row number is controlled by the FASCIATED EAR2 locus.** *Nat Genet* 2013, **45**:334-337.
- The authors mapped a maize kernel row number QTL near *FEA2*, the *CLV2* ortholog. They also used TILLING to generate weak *fea2* alleles and demonstrated that the natural variation in *FEA2* is important in controlling inflorescence meristem size and kernel row number.
44. Liu L, Du Y, Shen X, Li M, Sun W, Huang J, Liu Z, Tao Y, Zheng Y, Yan J *et al.*: **KRN4 controls quantitative variation in maize kernel row number.** *PLoS Genet* 2015, **11**:1-19.
- The authors fine-mapped a major kernel row number QTL, KRN4, which can enhance the productivity of maize. A ~1.2 kb transposon-containing insertion ~60 kb downstream of the *UB3* gene is likely responsible for the increase of KRN through affecting *UB3* expression and IM size. This study provides an example that variation in intergenic regions can regulate quantitative variation of important agronomic traits.
45. Zhang D, Sun W, Singh R, Zheng Y, Cao Z, Li M, Lunde C, Hake S, Zhang Z: **GRF-interacting factor1 (gif1) regulates shoot**

- architecture and meristem determinacy in maize. *Plant Cell* 2018, **30**:360-374.**
- The authors identified *gif1* mutants, which show an overproliferation of meristematic cells, resulting in a fasciated IM. The GIF1 protein regulates IM proliferation partially by targeting the *UB3* gene.
46. Du Y, Liu L, Li M, Fang S, Shen X, Chu J, Zhang Z: **UNBRANCHED3 regulates branching by modulating cytokinin biosynthesis and signaling in maize and rice.** *New Phytol* 2017, **214**:721-733.
 47. Pautler M, Eveland AL, LaRue T, Yang F, Weeks R, Lunde C, Je BI, Meeley R, Komatsu M, Vollbrecht E *et al.*: **FASCIATED EAR4 encodes a bZIP transcription factor that regulates shoot meristem size in maize.** *Plant Cell* 2015, **27**:104-120.
- The authors characterized a bZIP transcription factor FEA4 in meristem regulation, and found that it functions in parallel to the CLV-WUS pathway. FEA4 functions in the meristem periphery to promote differentiation by regulating auxin-based responses and leaf differentiation. *Note*: The *fea4* IM picture used in Table 1 is copyrighted by the American Society of Plant Biologists and is reprinted with permission.
48. Running MP, Meyerowitz EM: **Mutations in the PERIANTHIA gene of Arabidopsis specifically alter floral organ number and initiation pattern.** *Development* 1996, **122**:1261-1269.
 49. Yang F, Bui HT, Pautler M, Llaca V, Johnston R, Lee B-h, Kolbe A, Sakai H, Jackson D: **A maize glutaredoxin gene, Abphy12, regulates shoot meristem size and phyllotaxy.** *Plant Cell Online* 2015, **27**:121-131.
 50. Zeng J, Dong ZC, Wu HJ, Tian ZX, Zhao Z: **Redox regulation of plant stem cell fate.** *Embo J* 2017, **36**:2844-2855.
 51. Gallavotti A, Barazesh S, Malcomber S, Hall D, Jackson D, Schmidt RJ, McSteen P: **sparse inflorescence1 encodes a monocot-specific YUCCA-like gene required for vegetative and reproductive development in maize.** *Proc Natl Acad Sci U S A* 2008, **105**:15196-15201.
 52. Phillips KA, Skirpan AL, Liu X, Christensen A, Slewinski TL, Hudson C, Barazesh S, Cohen JD, Malcomber S, McSteen P: **vanishing tassels2 encodes a grass-specific tryptophan aminotransferase required for vegetative and reproductive development in maize.** *Plant Cell* 2011, **23**:550-566.
 53. Mashiguchi K, Tanaka K, Sakai T, Sugawara S, Kawaide H, Natsume M, Hanada A, Yaeno T, Shirasu K, Yao H *et al.*: **The main auxin biosynthesis pathway in Arabidopsis.** *Proc Natl Acad Sci U S A* 2011, **108**:18512-18517.
 54. Salehin M, Bagchi R, Estelle M: **SCFTIR1/AFB-based auxin perception: mechanism and role in plant growth and development.** *Plant Cell* 2015, **27**:9-19.
 55. Galli M, Liu Q, Moss BL, Malcomber S, Li W, Gaines C, Federici S, Roshkovan J, Meeley R, Nemhauser JL *et al.*: **Auxin signaling modules regulate maize inflorescence architecture.** *Proc Natl Acad Sci U S A* 2015, **112**:13372-13377.
 56. McSteen P, Malcomber S, Skirpan A, Lunde C, Wu XT, Kellogg E, Hake S: **barren inflorescence2 encodes a co-ortholog of the PINOID serine/threonine kinase and is required for organogenesis during inflorescence and vegetative development in maize.** *Plant Physiol* 2007, **144**:1000-1011.
 57. Skirpan A, Culler AH, Gallavotti A, Jackson D, Cohen JD, McSteen P: **BARREN INFLORESCENCE2 interaction with ZmPIN1a suggests a role in auxin transport during maize inflorescence development.** *Plant Cell Physiol* 2009, **50**:652-657.
 58. Huang P, Jiang H, Zhu CM, Barry K, Jenkins J, Sandor L, Schmutz J, Box MS, Kellogg EA, Brutnell TP: **Sparse panicle1 is required for inflorescence development in Setaria viridis and maize.** *Nat Plants* 2017, **3**.
 59. Leiboff S, Li X, Hu HC, Todt N, Yang J, Li X, Yu X, Muehlbauer GJ, Timmermans MCP, Yu J *et al.*: **Genetic control of morphometric diversity in the maize shoot apical meristem.** *Nat Commun* 2015, **6**:1-10.
- The authors captured SAM size variation within a diverse maize inbred panel by high-throughput image processing and demonstrated correlations between seedling SAM size and agronomic traits. The authors also used GWAS to identify SNPs and candidate genes that contribute to SAM size regulation.
60. Giullini A, Wang J, Jackson D: **Control of phyllotaxy by the cytokinin-inducible response regulator homologue ABPHYL1.** *Nature* 2004, **430**:1031-1034.
 61. Lee B-h, Johnston R, Yang Y, Gallavotti A, Kojima M, Travencolo BAN, Costa LdF, Sakakibara H, Jackson D: **Studies of aberrant phyllotaxy1 mutants of maize indicate complex interactions between auxin and cytokinin signaling in the shoot apical meristem.** *Plant Physiol* 2009, **150**:205-216.
 62. Woodward JB, Abeydeera ND, Paul D, Phillips K, Rapala-Kozik M, Freeling M, Begley TP, Ealick SE, McSteen P, Scanlon MJ: **A maize thiamine auxotroph is defective in shoot meristem maintenance.** *Plant Cell* 2010, **22**:3305-3317.
 63. Leonard A, Holloway B, Guo M, Rupe M, Yu G, Beatty M, Zastrow-Hayes G, Meeley R, Llaca V, Butler K *et al.*: **Tassel-less1 encodes a boron channel protein required for inflorescence development in maize.** *Plant Cell Physiol* 2014, **55**:1044-1054.
 64. Rosa M, Abraham-Juarez MJ, Lewis MW, Fonseca JP, Tian W, Ramirez V, Luan S, Pauly M, Hake S: **The maize MID-COMPLEMENTING ACTIVITY homolog CELL NUMBER REGULATOR13/NARROW ODD DWARF coordinates organ growth and tissue patterning.** *Plant Cell* 2017, **29**:474-490.
 65. Thompson BE, Basham C, Hammond R, Ding QY, Kakrana A, Lee TF, Simon SA, Meeley R, Meyers BC, Hake S: **The dicer-like1 homolog fuzzy tassel is required for the regulation of meristem determinacy in the inflorescence and vegetative growth in maize.** *Plant Cell* 2014, **26**:4702-4717.
 66. Chuck G, Whipple C, Jackson D, Hake S: **The maize SBP-box transcription factor encoded by tasselsheath4 regulates bract development and the establishment of meristem boundaries.** *Development* 2010, **137**:1585.
 67. Chuck G, Cigan AM, Saeteurn K, Hake S: **The heterochronic maize mutant Corngrass1 results from overexpression of a tandem microRNA.** *Nat Genet* 2007, **39**:544-549.
 68. Wu G, Poethig RS: **Temporal regulation of shoot development in Arabidopsis thaliana by miR156 and its target SPL3.** *Development* 2006, **133**:3539-3547.
 69. Doebley J, Stec A, Hubbard L: **The evolution of apical dominance in maize.** *Nature* 1997, **386**:485-488.
 70. Galinat WC: **The origin of maize.** *Annu Rev Genet* 1971, **5**:447-478.
 71. Doebley J: **The genetics of maize evolution.** *Annu Rev Genet* 2004, **38**:37-59.
 72. Studer A, Zhao Q, Ross-Ibarra J, Doebley J: **Identification of a functional transposon insertion in the maize domestication gene tb1.** *Nat Genet* 2011, **43**:1160-U1164.
 73. Whipple CJ, Kebrom TH, Weber AL, Yang F, Hall D, Meeley R, Schmidt R, Doebley J, Brutnell TP, Jackson DP: **grassy tillers1 promotes apical dominance in maize and responds to shade signals in the grasses.** *Proc Natl Acad Sci U S A* 2011, **108**:E506-E512.
 74. Dong ZB, Li W, Unger-Wallace E, Yang JL, Vollbrecht E, Chuck G: **Ideal crop plant architecture is mediated by tassels replace upper ears1, a BTB/POZ ankyrin repeat gene directly targeted by TEOSINTE BRANCHED1.** *Proc Natl Acad Sci U S A* 2017, **114**:E8656-E8664.
- The authors demonstrated *TRU1* is a direct target of TB1 and is overexpressed in modern maize, providing a molecular mechanism for how TB1 regulates axillary meristems and transformed teosinte into maize.
75. Munos S, Ranc N, Botton E, Berard A, Rolland S, Duffe P, Carretero Y, Le Paslier MC, Delalande C, Bouzayen M *et al.*: **Increase in tomato locule number is controlled by two single-nucleotide polymorphisms located near WUSCHEL.** *Plant Physiol* 2011, **156**:2244-2254.
 76. Xu C, Liberatore KL, MacAlister CA, Huang ZJ, Chu YH, Jiang K, Brooks C, Ogawa-Ohnishi M, Xiong GY, Pauly M *et al.*: **A cascade**

of arabinosyltransferases controls shoot meristem size in tomato. *Nat Genet* 2015, **47**:784-792.

77. Rodriguez-Leal D, Lemmon ZH, Man J, Bartlett ME, Lippman ZB:
•• **Engineering quantitative trait variation for crop improvement by genome editing.** *Cell* 2017, **171**:470-480.

The authors show that CRISPR/Cas9 genome editing of promoters generates diverse cis-regulatory alleles that improve the yield of tomato. By segregating away the CRISPR/Cas9 machinery in the following generation, they fixed a wide range of stable promoter alleles, which can be used in breeding.