



Maturation-dependent vulnerability of emotion regulation as a response to COVID-19 related stress in adolescents

Patrícia Gerván^{a,b,*}, Nóra Bunford^c, Katinka Utczás^d, Zsófia Tróznai^d, Gyöngyi Oláh^{b,e,f}, Hanna Szakács^{e,f}, Pálma Kriston^g, Ferenc Gombos^{b,f}, Ilona Kovács^{b,c,f}

^a Institute of Psychology, Pázmány Péter Catholic University, Budapest 1088, Hungary

^b Adolescent Development Research Group, Hungarian Academy of Sciences - Pázmány Péter Catholic University, Budapest 1088, Hungary

^c Institute of Cognitive Neuroscience and Psychology, Res. Centre for Natural Sciences, Budapest 1117, Hungary

^d Research Centre for Sport Physiology, University of Physical Education, Budapest 1123, Hungary

^e Doctoral School of Mental Health Sciences, Semmelweis University, Budapest 1089, Hungary

^f Laboratory for Psychological Research, Pázmány Péter Catholic University, Budapest 1088, Hungary

^g Doctoral School of Education University of Szeged, Faculty of Humanities and Social Sciences, 6722, Hungary

ARTICLE INFO

Article history:

Received 19 January 2022

Revised 16 August 2022

Accepted 20 August 2022

Available online xxxx

Keywords:

COVID-19

Adolescence

Emotion regulation

DERS

Pubertal maturation

Bone age

ABSTRACT

Background: The COVID-19 pandemic created unpredictable circumstances resulting in increased psychological strain. Here we investigate pandemic-related alterations in emotion regulation in adolescents assessed before and during the pandemic. We also take biological age into account in the response to the pandemic.

Methods: Mann-Whitney *U* tests were conducted to compare baseline data on the Difficulties in Emotion Regulation Scale (DERS) total scores of a pre-pandemic adolescent cohort ($n = 241$) with those obtained during the second wave of the pandemic ($n = 266$). We estimated biological age based on an ultrasonic boneage assessment procedure in a subgroup of males, including grammar school and vocational school students in the 9th and 10th grades, and analyzed their data independently.

Findings: There is a gender difference in the timing of vulnerability for pandemic-related stress in grammar school students: females are affected a year earlier than males. Vocational school male students mature faster than grammar school male students, and the timing of emotional vulnerability also precedes that of the grammar school students'.

Discussion: We interpret our findings within a developmental model suggesting that there might be a window of highest vulnerability in adolescent emotion regulation. The timing of the window is determined by both chronological and biological age, and it is different for females and males.

Application to practice: Defining the exact temporal windows of vulnerability for different adolescent cohorts allows for the timely integration of preventive actions into adolescent care to protect mental health during future chronic stressful situations.

© 2022 Published by Elsevier Inc.

Introduction

Besides the health and life threatening consequences, Covid-19 pandemic resulted in psychological distress and disturbance in the population, including children and adolescents. Studies on the psychological consequences of COVID-19 revealed an elevation in the perceived stress by adolescents (Mohler-Kuo et al., 2021; Nocentini et al., 2021; Zhang et al., 2020). Adolescents have been facing multiple adversaries since

the start of the pandemic: with schools closed, their daily routines are disrupted; lack of personal contact with peers and other containment measures increase the prevalence of depressive and anxiety symptoms (Ellis et al., 2020; Zhou et al., 2020); attending school in-person once the lockdown is lifted is a challenge for some students, especially for those with mental health problems, such as depression (Lee, 2020). Effectively managing and attenuating emotions helps prevent being overwhelmed by negative emotions and promotes daily functioning of adolescents in challenging life situations (Bunford et al., 2015; Compas et al., 2017). Accordingly, under these extreme circumstances, the acquisition of age-appropriate emotion regulatory skills becomes one of the most important developmental tasks, and the maturity of the

* Corresponding author at: Institute of Psychology, Pázmány Péter Catholic University, 1 Mikszáth sq., Budapest 1088, Hungary.

E-mail address: gervan.patricia@btk.ppke.hu (P. Gerván).

emotion processing brain circuitry in the background seems to be a particularly relevant aspect of healthy functioning (Ely et al., 2021; Zubovics et al., 2021).

Under regular circumstances, by the time youth reach adolescence, they have acquired differentiated and diversified emotion expression and regulation skills and are able to incorporate environmental feedback and accordingly modify their emotional expressions and regulatory responses (Bunford & Evans, 2017). On the other hand, completely adult-like emotion regulation cannot be expected since the biological changes of adolescence involve brain structures and systems implicated in emotion expression and regulation as well. There is, for example, a second surge of synaptogenesis (dendritic pruning and myelinogenesis), making the teenage years one of the most dynamic periods of human development (Arain et al., 2013; Lynch et al., 2020). Due to the considerable increases in sex hormones, neurocircuitry is still functionally and structurally unstable resulting in marked vulnerability (Patel et al., 2021). Extensive maturation of the basal ganglia and frontal lobe occurs during adolescence (Larsen & Luna, 2015; Sowell et al., 2004), with both the basal ganglia (Johnson et al., 2003) and the frontolimbic system (Banks et al., 2007) implicated in emotion generation and regulation. The prefrontal cortex remains under construction and there is a decrease in dopamine and serotonin levels ((Chugani et al., 1999; Wahlstrom et al., 2010), with these neurotransmitters also involved in emotion regulation (GABA: Thayer & Lane, 2000; dopamine: Laviolette, 2007; serotonin transporter gene: Canli & Lesch, 2007).

Moderated by age and context, there are gender differences in emotion regulation. Females employ both more adaptive (e.g., active coping and re-evaluation) and maladaptive (e.g., rumination and suppression) emotion regulatory strategies relative to males (Chaplin & Aldao, 2013), with this difference potentially reflecting the more general tendency of women being more aware of their emotions and more open to engaging with their emotions (Nolen-Hoeksema et al., 1999). Females tend to show better emotion regulation in young adolescence (age 9–12 years) but worse emotion regulation than males in middle adolescence (age 13–16 years, Zimmer-Gembeck & Skinner, 2011), highlighting a stage of increased emotional vulnerability for females during the middle adolescent years as also reflected in the frequency of anxiety-depressive problems during this developmental period (Muris et al., 2000).

There is a relative paucity of empirical research on the direct effect of acute, laboratory-induced stress on emotion regulation in children and adolescents (Langer et al., 2020), and the available data are mixed, with some evidence indicating that acute stress impairs emotion regulation (Raio et al., 2013; Raio & Phelps, 2015) and other findings suggesting improvements in emotion regulation (Kinner et al., 2014; Langer et al., 2020). Regarding the effects of acute, non-laboratory induced stress, findings on child and adolescent survivors of disasters, wars, and other acute events are also inconsistent regarding behavioral and emotional outcomes. Some data indicate an increase in behavioral (e.g., aggression) and emotional (e.g., anxiety, depression) problems (Marsee, 2008; Scott, Lapré, Marsee, & Weems, 2014) whereas others show no adverse effects on these outcomes (Durkin et al., 1993).

We would like to investigate the impact of Covid-19 pandemic events on emotion regulation, and to determine the modulatory effects of gender and/or maturational status on the timing and magnitude of the pandemic impact. In addition to assuming a gender difference in the timing of the strongest impact, we were also interested in those potential windows of vulnerability that are suggested by the above-mentioned brain maturational processes (e.g., pruning, late frontolimbic maturation, neurotransmitter imbalance). We applying a chronological definition of adolescence which refers to the time between the beginning of puberty and adulthood. Considering the large individual differences in the onset of puberty, to ensure that most of our participants reach adolescence and be above 14 years of age (Farello et al., 2019), we focused on a cohort of students in the 9th and 10th grades. We relied on grades instead of chronological age to assure that their previous

cognitive and social experience was similar. In addition to age, we were also interested in contextual effects that might have an impact on maturation, therefore, we included both grammar school and vocational school students. On a subgroup of students, we also employed bone age measurements to assess their biological maturity levels and to see how biological age might be associated with emotion regulation development. A unique characteristic of our study is that we compare pre-pandemic and pandemic emotion regulation results within age-groups and genders.

Methods

Participants

To study the impact of the COVID-19 pandemic on emotion regulation of adolescents, we compared a Pre-Pandemic and a Pandemic cohort. Both cohorts included students from different grammar schools (GS) attending 9th or 10th grade and a vocational school (VS) attending 9th grade. In Hungary, secondary education includes three traditional types of schools focused on different academic levels: grammar schools, secondary vocational schools, and technical schools. Grammar schools offer general academic education and provides the students with a general qualification for higher education. Vocational schools offer vocational education as well as practical training and students acquire a qualification at the end of their studies. Descriptive statistics of the Pre-Pandemic and Pandemic cohorts are summarized in Table 1. in the Results section.

We involved 170 participants in the Pre-Pandemic GS groups, and 208 participants in the Pandemic GS groups, 71 participants in the Pre-Pandemic VS group, and 58 participants in the Pandemic VS group (see details in Table 1.). (The grammar school participants were from six different institutions. The average number of pupils in these schools is 675, and an average of 107 pupils attend 9th, and 86 pupils attend 10th grade.)

VS participants are from BGSZC Bánki Donát Vocational High School of Transport Engineering (total number of pupils is 840, 124 and 107 students attend 9th and 10th grades, respectively). The proportion of female students is <1% in this school. Due to the low proportion of females all our VS participants are males. It is important to note that VS and GS students have significant differences in economic, social and cultural background. VS students have a typically lower socio-economic background.

127 9th grader male students took part in bone age assessments (see details in Table 2.).

All of our subjects were Caucasian.

Tools

Questionnaire

Emotion regulation processes were measured by the validated Hungarian version (Kökönyei et al., 2014; Cronbach's alpha = 0.80) of the Difficulties in Emotion Regulation Scale (DERS; Gratz & Roemer, 2004; Cronbach's alphas > 0.80). The DERS is a comprehensive, 36-item, self-report measure that assesses the typical levels of emotion dysregulation overall and across the following dimensions: (1) nonacceptance of emotional responses, (2) difficulties engaging in goal-directed behaviors when distressed, (3) difficulties controlling impulsive behaviors, (4) lack of emotional awareness, (5) limited access to emotion regulation strategies perceived as effective, (6) lack of emotional clarity. The dimensions of DERS seem to correlate with psychological problems reflecting emotion dysregulation, specifically depression, anxiety, suicidal ideation, eating disorders, alcohol and drug use, ADHD, and social impairment (Bunford et al., 2015, 2018; Weinberg & Klonsky, 2009). It has been validated on adolescents (11–17 years of age) and had a good internal consistency (Bunford et al., 2015; Neumann et al., 2010).

Table 1

Mean ages and DERS results of the subgroups. Mann-Whitney *U* test results of DERS total scores for Pre-pandemic and Pandemic subgroups. Gr = Grade, Mdn = median, GS = Grammar School, VS=Vocational School (**p* < 0.05 > ***p* < 0.01).

Group	Gr	Subgroup	N	Mean age (in years)	SD	Mean DERS total	SD	Mdn DERS total	Mean Rank	Sum of Ranks	U	P
GS female	9th	Pre-Pandemic	63	14.82	0.49	80.39	19.48	78	53.98	3401	1385	0.01*
		Pandemic	59	15.16	0.35	90.76	24.18	85	69.53	4102		
	10th	Pre-Pandemic	58	15.76	0.43	83.58	22.12	81.5	56.54	3279	1505	0.85
		Pandemic	53	16.16	0.44	83.54	24.88	80	55.41	2936		
GS male	9th	Pre-Pandemic	22	14.94	0.65	80.00	14.61	80.5	39.00	858	561	0.79
		Pandemic	53	15.39	0.42	79.35	16.52	76	37.58	1992		
	10th	Pre-Pandemic	27	15.96	0.43	72.96	17.50	70	28.83	778	400	0.03*
		Pandemic	43	16.55	0.45	84.41	23.07	80	39.69	1706		
VS male	9th	Pre-Pandemic	71	15.49	0.55	67.40	17.10	64	49.40	3507	951	0.00**
		Pandemic	58	15.47	0.45	89.41	25.50	84.5	84.09	4877		

Bone age assessment

Skeletal maturity (bone age) was assessed with an ultrasonic device (Sunlight BoneAge, Sunlight Medical Ltd., Tel Aviv, Israel). Bone age measurements were performed on the left hand and wrist region of the subjects. The same experimenter performed all the bone age assessments with the same device. The device estimates bone age (in years and months) by measuring the speed of sound and the distance between the transducers, using algorithms based on gender and ethnicity (see the detailed description of the method in Kovács et al., 2022).

Procedure

Administering the DERS: Pre-pandemic dataset

The Pre-pandemic GS dataset, which includes both female and male participants, was derived from two databases of different Hungarian adolescent research projects. The first database was completed in 2016 (a previous analysis on these data, not related to the current manuscript, has been published in Kriston & Pikó, 2018). The questionnaires were administered in a paper-and-pencil format in schools.

The other database, provided by the Adolescent Developmental and Translational Neuroscience Research Group, contains data from 2019 and early 2020 (January and February), from both male and female participants. Data collection was administered in the research center via the Qualtrics software, Version 2020 (Qualtrics, Provo, UT).

VS males have been participating in a longitudinal investigation of the Adolescent Development Research Group supported by the Hungarian Academy of Sciences and Pázmány Péter Catholic University. We recruited participants in collaboration with the school management, and participants were given credits in their school for the participation. The Pre-Pandemic assessment took place in November 2019. Students completed the questionnaire in the computer lab of their school using the PsyToolkit software (Stoet, 2010, 2017).

Administering the DERS: Pandemic dataset

Data collection for the Pandemic GS cohort was carried out between November 2020 and February 2021, during the second wave of COVID-19 in Hungary, when the levels of new cases and COVID-related deaths were among the highest in Europe. During this time, Hungary implemented strict pandemic-related restrictions, which included national quarantine with the closure of schools and shops (except for daily essentials), movie theatres, restaurants, and a curfew after 8 p.m. All gatherings were forbidden, private and family events could be held for up to

Table 2

Means and standard deviations of chronological and bone age for the two male cohorts.

Group	N	Chron. age (mean in y)	SD	Bone age (mean in y)	SD
GS male 9 th Grade	31	15.33	0.46	15.50	1.49
VS male 9 th Grade	96	15.40	0.57	16.08	1.19

10 people. In a convenience sampling procedure, GS students were invited to join the study via an invitation e-mail sent out by either the school psychologist or the teachers. We asked 9th and 10th grade students to simply fill out the online questionnaire, and we did not apply any exclusion criteria.

GS students were offered extra accountable hours in their volunteer work projects, which are mandatory in Hungarian secondary schools, for participating in the research. Due to the epidemiological situation and the lockdown, the data were gathered online via the Qualtrics software.

The VS male students in the Pandemic subgroup participated between November 2020 and February 2021. Similarly to the GS subjects, the data were gathered online using Qualtrics.

Assessing skeletal maturation

Ultrasonic bone age estimations were carried out before (November 2020) and after the lockdowns (April 2021). The procedure took place either at the respective schools of the participants or at the Research Center for Sport Physiology at the University of Physical Education, Budapest.

Ethics

The PPCU Institute of Psychology Committee for Research in Psychology (reference number 2020_30) and the Hungarian United Ethical Review Committee for Research in Psychology (reference number 2017/84) approved the study. The pre-pandemic grammar school data collection by the Adolescent Developmental and Translational Neuroscience Research Group was approved by the National Institute of Pharmacy and Nutrition (OGYÉI/17089–8/2019).

Written informed consent was obtained from all subjects and their parents.

Results

Pre-Pandemic vs. Pandemic DERS scores

Mean DERS total score of subgroups is shown in Fig. 1. We compared Pre-pandemic and Pandemic DERS total scores in the groups of 9th and 10th graders. The distribution of total DERS scores was abnormal (Shapiro–Wilk normality test >0.00). Mann-Whitney *U* tests were conducted to determine if the differences between the Pre-Pandemic and Pandemic subgroups regarding the DERS total score were significant. DERS results are summarized in Table 1.

We found a significantly higher DERS median score in the subgroup of 9th grader Pandemic GS females than in the Pre-Pandemic subgroup. In contrast, we did not find a significant difference between the Pandemic and Pre-Pandemic subgroups of GS males. In the 9th grader VS male group, the median of the Pandemic subgroup was higher as




Fig. 1. DERS total score means of the Pre-Pandemic (empty symbols) and Pandemic (solid symbols) subgroups. Error bars indicate 2SE. Asterisks indicate significant difference in the Mann-Whitney U test (* $p < 0.05$; ** $p < 0.01$). In the Grammar School (GS) female population, the 9th grader Pandemic subgroup has higher DERS total scores as compared to Pre-Pandemic, while there is no difference between the two subgroups in 10th grade. This pattern of difference was the reverse in the GS male population: there is no significant difference between the Pre-Pandemic and Pandemic subgroups in 9th grade, whereas a significant difference occurred between the 10th grader subgroups. The difference between the subgroups of the 9th grader Vocational School (VS) males was similar to the 9th grader GS female pattern: the Pandemic subgroup showed significantly higher DERS total scores than the Pre-Pandemic.

compared to that of the Pre-Pandemic subgroup (similarly to the 9th grader GS female group).

The reverse of the above-described pattern occurred in the groups of 10th graders. The Pandemic subgroup of GS females did not show a significant change in the medians of the total DERS scores as compared to that of the Pre-pandemic subgroup. In contrast, the 10th grader GS male Pandemic subgroup showed a significant increase in the total DERS score median as compared to that of the Pre-Pandemic subgroup.

To compare the GS and VS males on chronological and bone age variables, a univariate analysis of variance (ANOVA) was carried out. ANOVA revealed no difference between samples on chronological age ($F(1,125) = 0.38, p = 0.54$). Conversely, bone age of VS males was significantly higher than that of GS males ($F(1,125) = 4.81, p = 0.03$). We ran paired sample *t*-tests to determine whether chronological age and bone age is dissociated within these two groups of males. We did not find a significant difference in these variables in GS males ($t(30) = -0.675, p = 0.505$). In contrast, there was a substantial difference in the VS male group between chronological and bone age ($t(95) = -5.29, p < 0.00$), bone age being higher by an average of 0.67 years (cc. 8 months) than chronological age.

Discussion

Interestingly, while emotion regulation (as assessed by the total score on DERS) in 9th grader GS females of the Pandemic cohort seemed to be significantly above that of the Pre-Pandemic subgroup (indicating more regulation problems), 10th grader Pre-Pandemic females provided very similar scores as those in the Pandemic group. Males going to GS had the opposite pattern: 9th graders were not affected by the Pandemic in emotion regulation according to their DERS scores, while those in the 10th grade Pandemic group scored significantly above (indicating more emotion regulation problems) the Pre-Pandemic group. Gender difference in the timing of puberty is an acknowledged fact, females typically outstrip males by one and a half years (Farello et al., 2019; Hoyt et al., 2020). This pattern of results, therefore, not only points to a gender difference in the timing of vulnerability but indicates a “window” of vulnerability in both genders: females being very sensitive to the stress due to the pandemic circumstances in the 9th grade, and more stable

after that; males being unaffected in the 9th grade, and vulnerable a year later.

We were also particularly interested in those potential windows of vulnerability that are suggested by the earlier mentioned brain maturational processes, e.g., cortical pruning, late frontolimbic maturation (Hwang et al., 2016; Marek et al., 2018; Yang & Tseng, 2021) and neurotransmitter imbalance (for a review see Pitzer, 2019). To this end, we assessed the biological age of both GS and VS adolescent males assuming that biological age as assessed by bone age will be in correlation with brain maturation (Kovács et al., 2022). We found that VS males are significantly more mature than GS males. This is in good agreement with a recent finding by Oelkers et al. (2020) who demonstrated that earlier pubertal onset is more likely in the group of lower socioeconomic status schoolboys as compared to peers with higher socioeconomic status.

9th grader VS males showed a decline in emotional regulation during the pandemic similarly to GS females, therefore, these earlier maturing males show an earlier window of vulnerability as compared to GS males.

As in the case of any developmental event, changes of emotion regulation skills will reach the point of fastest change in between the levels of childlike and adultlike emotion regulation. Since we are not assuming stepwise development, a sigmoid function is a good approximation of that, and the inflection point of this curve characterizes the point of fastest change (Burchinal & Appelbaum, 1991). The main idea of our hypothesis is that around the point of fastest change, there is a window of heightened vulnerability as well (Charmandari et al., 2003; Larsen & Luna, 2018; Semple et al., 2013). As we detailed in the introduction, emotion regulation is dependent on the maturation of brain structures and systems as well as on age, gender, and contextual factors. These curves, drawn hypothetically in Fig. 2, for the three cohorts that we studied, very clearly show the relevance of a multifactorial view of development that is not simply dependent on chronological age.

Practice implications

The practical message of our results combined with the theoretical scenario described in Fig. 2, is that adolescence is not simply a period of increased vulnerability in the emotional domain, but it is a




Fig. 2. Schematic representation of the adolescent development of emotional regulation. The hypothetical functions each represent the adolescent course of development between the end of childhood and adulthood. The fastest developmental change occurs at the inflection points of these curves. Assuming similar developmental speed (slope of the curve) in earlier and later maturing groups, the inflection points define the centers of windows of highest vulnerability where development and plasticity are fastest, and vulnerability is greatest. (GS = grammar school, VS = vocational school.)

fine-grained developmental process, with potentially narrow windows for heightened sensitivity and vulnerability. A number of psychopathologies first emerge or manifest in adolescence, or show a dramatic increase in terms of prevalence during adolescence (Giedd et al., 2008). More accurate exploration of the precise timing of these vulnerability windows across the adolescent population would allow health and education professionals to properly time interventions and help in the most critical periods to achieve the best impact. For example, our results suggest that while Caucasian GS females and VS males may need the most intensive attention and help in the 9th grade, Caucasian GS males will need it a grade later. However, we emphasize that the individual variability in timing is very extensive, there are large individual differences in the onset and speed of puberty (Dorn & Biro, 2011; Farello et al., 2019). Alterations in the timing of puberty with respect to peers of the same age has a potential risk of receiving less attention and help at the right time. Several potentially detrimental impacts of early (Mendle et al., 2007; Mendle & Ferrero, 2012) and late puberty (Graber, 2013; Negriff & Susman, 2011) have been identified in the literature, and lifelong negative impacts of these conditions have been demonstrated as well (Graber et al., 2004; Natsuaki et al., 2009). We have shown that the timing of pubertal maturation is related to the window of heightened vulnerability in emotional regulation, therefore intensive attention in pediatric care should be given accordingly, taking the altered maturational speed into account.

Limitations

Finally, a number of potential limitations need to be considered. The most relevant constraint is that exclusively Caucasian students participated in the current study, and the findings on the timing of vulnerability windows may not be transferable to other ethnic groups (see e.g. Herman-Giddens, 2006; Seaton & Carter, 2019). Our study was also limited in age-range. By expanding the age-range, it might be interesting to see whether the one-year vulnerability window extends to earlier or later grades. We hope, on the other hand, that the COVID-19 pandemic will not provide the grounds for such an investigation because - with the vaccinations introduced widely - the need for constraining the youth will subside.

Conclusion

According to our original purpose, we were able to assess the immediate impact of the current pandemic events on emotion regulation in

adolescents. We found a gender difference in the timing of the strongest negative impact; females being affected about a year earlier than males in the GS population. However, it seems that earlier VS maturing males are closer in terms of emotional maturity to GS females than to GS males. This is a novel and very surprising finding as similar comparisons have not been made before. The results might corroborate our hypothesis with respect to the maturational effects in emotional regulation that should determine when a particular individual is most sensitive to stressful events.

Acknowledgments

This work was supported by the National Research, Development and Innovation Office of Hungary (Grant K-134370, 2020) to I.K., and by the Eötvös Loránd Research Network, Hungary (ELRN-PPCU Adolescent Development Research Group).

We would like to thank Bettina Pikó for her contribution.

References

- Arain, M., Haque, M., Johal, L., Mathur, P., Nel, W., Rais, A., Sandhu, R., & Sharma, S. (2013). Maturation of the adolescent brain. *Neuropsychiatric Disease and Treatment*, 9, 449–461. PubMed <https://doi.org/10.2147/NDT.S39776>.
- Banks, S. J., Eddy, K. T., Angstadt, M., Nathan, P. J., & Phan, K. L. (2007). Amygdala–frontal connectivity during emotion regulation. *Social Cognitive and Affective Neuroscience*, 2(4), 303–312. <https://doi.org/10.1093/scan/nsm029>.
- Bunford, N., & Evans, S. W. (2017). Emotion regulation and social functioning in adolescence: Conceptualization and treatment. *School mental health services for adolescents* (pp. 161–181) (1st ed.). Oxford University Press. <https://doi.org/10.1093/med-psych/9780199352517.003.0008>.
- Bunford, N., Evans, S. W., Becker, S. P., & Langberg, J. M. (2015). Attention-deficit/hyperactivity disorder and social skills in youth: A moderated mediation model of emotion dysregulation and depression. *Journal of Abnormal Child Psychology*, 43(2), 283–296. PubMed. <https://doi.org/10.1007/s10802-014-9909-2>.
- Bunford, N., Evans, S. W., & Langberg, J. M. (2018). Emotion dysregulation is associated with social impairment among young adolescents with ADHD. *Journal of Attention Disorders*, 22(1), 66–82. <https://doi.org/10.1177/1087054714527793>.
- Burchinal, M., & Appelbaum, M. I. (1991). Estimating individual developmental functions: Methods and their assumptions. *Child Development*, 62(1), 23–43. JSTOR. <https://doi.org/10.2307/1130702>.
- Canli, T., & Lesch, K. -P. (2007). Long story short: The serotonin transporter in emotion regulation and social cognition. *Nature Neuroscience*, 10(9), 1103–1109. <https://doi.org/10.1038/nn1964>.
- Chaplin, T. M., & Aldao, A. (2013). Gender differences in emotion expression in children: A meta-analytic review. *Psychological Bulletin*, 139(4), 735–765. PubMed. <https://doi.org/10.1037/a0030737>.
- Charmandari, E., Kino, T., Souvatzoglou, E., & Chrousos, G. P. (2003). Pediatric stress: Hormonal mediators and human development. *Hormone Research in Paediatrics*, 59(4), 161–179. <https://doi.org/10.1159/000069325>.

- Chugani, D. C., Muzik, O., Behen, M., Rothermel, R., Janisse, J. J., Lee, J., & Chugani, H. T. (1999). Developmental changes in brain serotonin synthesis capacity in autistic and nonautistic children. *Annals of Neurology*, 45(3), 287–295. [https://doi.org/10.1002/1531-8249\(199903\)45:3<287::AID-ANA3>3.0.CO;2-9](https://doi.org/10.1002/1531-8249(199903)45:3<287::AID-ANA3>3.0.CO;2-9).
- Compas, B. E., Jaser, S. S., Bettis, A. H., Watson, K. H., Gruhn, M. A., Dunbar, J. P., ... Thigpen, J. C. (2017). Coping, emotion regulation, and psychopathology in childhood and adolescence: A meta-analysis and narrative review. *Psychological Bulletin*, 143(9), 939–991. <https://doi.org/10.1037/bul0000110>.
- Dorn, L. D., & Biro, F. M. (2011). Puberty and its measurement: A decade in review. *Journal of Research on Adolescence*, 21(1), 180–195. <https://doi.org/10.1111/j.1532-7795.2010.00722.x>.
- Durkin, M. S., Khan, N., Davidson, L. L., Zaman, S. S., & Stein, Z. A. (1993). The effects of a natural disaster on child behavior: Evidence for posttraumatic stress. *American Journal of Public Health*, 83(11), 1549–1553. <https://doi.org/10.2105/AJPH.83.11.1549>.
- Ellis, W. E., Dumas, T. M., & Forbes, L. M. (2020). Physically isolated but socially connected: Psychological adjustment and stress among adolescents during the initial COVID-19 crisis. *Canadian Journal of Behavioural Science / Revue Canadienne Des Sciences Du Comportement*, 52(3), 177–187. <https://doi.org/10.1037/cbs0000215>.
- Ely, B. A., Liu, Q., DeWitt, S. J., Mehra, L. M., Alonzo, C. M., & Gabbay, V. (2021). Data-driven parcellation and graph theory analyses to study adolescent mood and anxiety symptoms. *Translational Psychiatry*, 11(1), 266. <https://doi.org/10.1038/s41398-021-01321-x>.
- Farello, G., Altieri, C., Cutini, M., Pozzobon, G., & Verrotti, A. (2019). Review of the literature on current changes in the timing of pubertal development and the incomplete forms of early puberty. *Frontiers in Pediatrics*, 7, 147. <https://doi.org/10.3389/fped.2019.00147>.
- Giedd, J. N., Keshavan, M., & Paus, T. (2008). Why do many psychiatric disorders emerge during adolescence? *Nature Reviews Neuroscience*, 9(12), 947–957. <https://doi.org/10.1038/nrn2513>.
- Graber, J. A. (2013). Pubertal timing and the development of psychopathology in adolescence and beyond. *Puberty and Adolescence*, 64(2), 262–269. <https://doi.org/10.1016/j.jybeh.2013.04.003>.
- Graber, J. A., Seeley, J. R., Brooks-Gunn, J., & Lewinsohn, P. M. (2004). Is pubertal timing associated with psychopathology in young adulthood. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43(6), 718–726. <https://doi.org/10.1097/01.chi.0000120022.14101.11>.
- Gratz, K. L., & Roemer, L. (2004). Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in emotion regulation scale. *Journal of Psychopathology and Behavioral Assessment*, 26(1), 41–54. <https://doi.org/10.1023/B:JOBA.0000007455.08539.94>.
- Herman-Giddens, M. E. (2006). Recent data on pubertal milestones in United States children: The secular trend toward earlier development. *International Journal of Andrology*, 29(1), 241–246. <https://doi.org/10.1111/j.1365-2605.2005.00575.x>.
- Hoyt, L. T., Niu, L., Pachucki, M. C., & Chaku, N. (2020). Timing of puberty in boys and girls: Implications for population health. *SSM - Population Health*, 10, 100549–100549. <https://doi.org/10.1016/j.ssmph.2020.100549>.
- Hwang, K., Ghuman, A. S., Manoach, D. S., Jones, S. R., & Luna, B. (2016). Frontal preparatory neural oscillations associated with cognitive control: A developmental study comparing young adults and adolescents. *NeuroImage*, 136, 139–148. <https://doi.org/10.1016/j.neuroimage.2016.05.017>.
- Johnson, P. A., Hurley, R. A., Benkelfat, C., Herpertz, S. C., & Taber, K. H. (2003). Emotion Regulation in Borderline Personality Disorder: Contributions of Neuroimaging. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 15(4), 397–402. <https://doi.org/10.1176/appi.neuropsych.15.4.397>.
- Kinner, V. L., Het, S., & Wolf, O. T. (2014). Emotion regulation: Exploring the impact of stress and sex. *Frontiers in Behavioral Neuroscience*, 8, 397. <https://doi.org/10.3389/fnbeh.2014.00397>.
- Kökönyei, G., Urbán, R., Reinhardt, M., Józán, A., & Demetrovics, Z. (2014). The difficulties in emotion regulation scale: Factor structure in chronic pain patients. *Journal of Clinical Psychology*, 70(6), 589–600. <https://doi.org/10.1002/jclp.22036>.
- Kovács, I., Kovács, K., Gerván, P., Utczás, K., Oláh, G., Tróznai, Z., Berencsi, A., Szakács, H., & Gombos, F. (2022). Ultrasonic bone age fractionates cognitive abilities in adolescence. *Scientific Reports*, 12(1), 5311. <https://doi.org/10.1038/s41598-022-09329-z>.
- Kriston, P., & Pikó, B. (2018). Érzelemszabályozás és figyelemkontroll középiskolások körében. *NEVELÉSTUDOMÁNY: OKTATÁS KUTATÁS INNOVÁCIÓ*, 6(4), 83–94. <https://doi.org/10.21549/NTNY.24.2018.4.7>.
- Langer, K., Hagedorn, B., Stock, L., Otto, T., Wolf, O. T., & Jentsch, V. L. (2020). Acute stress improves the effectiveness of cognitive emotion regulation in men. *Scientific Reports*, 10(1), 11571. <https://doi.org/10.1038/s41598-020-68137-5>.
- Larsen, B., & Luna, B. (2015). In vivo evidence of neurophysiological maturation of the human adolescent striatum. *Developmental Cognitive Neuroscience*, 12, 74–85. <https://doi.org/10.1016/j.dcn.2014.12.003>.
- Larsen, B., & Luna, B. (2018). Adolescence as a neurobiological critical period for the development of higher-order cognition. *Neuroscience & Biobehavioral Reviews*, 94, 179–195. <https://doi.org/10.1016/j.neubiorev.2018.09.005>.
- Lavolette, S. R. (2007). Dopamine modulation of emotional processing in cortical and subcortical neural circuits: Evidence for a final common pathway in schizophrenia? *Schizophrenia Bulletin*, 33(4), 971–981. <https://doi.org/10.1093/schbul/sbm048>.
- Lee, J. (2020). Mental health effects of school closures during COVID-19. *The Lancet Child & Adolescent Health*, 4(6), 421. [https://doi.org/10.1016/S2352-4642\(20\)30109-7](https://doi.org/10.1016/S2352-4642(20)30109-7).
- Lynch, K. M., Cabeen, R. P., Toga, A. W., & Clark, K. A. (2020). Magnitude and timing of major white matter tract maturation from infancy through adolescence with NODDI. *NeuroImage*, 212, 116672–116672. <https://doi.org/10.1016/j.neuroimage.2020.116672>.
- Marek, S., Tervo-Clemmens, B., Klein, N., Foran, W., Ghuman, A. S., & Luna, B. (2018). Adolescent development of cortical oscillations: Power, phase, and support of cognitive maturation. *PLoS Biology*, 16(11), e2004188–e2004188. <https://doi.org/10.1371/journal.pbio.2004188>.
- Marsee, M. A. (2008). Reactive aggression and posttraumatic stress in adolescents affected by hurricane Katrina. *Journal of Clinical Child & Adolescent Psychology*, 37(3), 519–529. <https://doi.org/10.1080/15374410802148152>.
- Mendle, J., & Ferrero, J. (2012). Detrimental psychological outcomes associated with pubertal timing in adolescent boys. *Developmental Review*, 32(1), 49–66. <https://doi.org/10.1016/j.dr.2011.11.001>.
- Mendle, J., Turkheimer, E., & Emery, R. E. (2007). Detrimental psychological outcomes associated with early pubertal timing in adolescent girls. *Developmental Review*, 27(2), 151–171. <https://doi.org/10.1016/j.dr.2006.11.001>.
- Mohler-Kuo, M., Dzemali, S., Foster, S., Werlen, L., & Walitza, S. (2021). Stress and mental health among children/adolescents, their parents, and young adults during the first COVID-19 lockdown in Switzerland. *International Journal of Environmental Research and Public Health*, 18(9). <https://doi.org/10.3390/ijerph18094668>.
- Muris, P., Schmidt, H., & Merckelbach, H. (2000). Correlations among two self-report questionnaires for measuring DSM-defined anxiety disorder symptoms in children: The screen for child anxiety related emotional disorders and the Spence Children's anxiety scale. *Personality and Individual Differences*, 28(2), 333–346. [https://doi.org/10.1016/S0191-8869\(99\)00102-6](https://doi.org/10.1016/S0191-8869(99)00102-6).
- Natsuaki, M. N., Biehl, M. C., & Ge, X. (2009). Trajectories of depressed mood from early adolescence to young adulthood: The effects of pubertal timing and adolescent dating. *Journal of Research on Adolescence*, 19(1), 47–74. <https://doi.org/10.1111/j.1532-7795.2009.00581.x>.
- Negriff, S., & Susman, E. J. (2011). Pubertal timing, depression, and externalizing problems: A framework, review, and examination of gender differences. *Journal of Research on Adolescence*, 21(3), 717–746. <https://doi.org/10.1111/j.1532-7795.2010.00708.x>.
- Neumann, A., van Lier, P. A. C., Gratz, K. L., & Koot, H. M. (2010). Multidimensional assessment of emotion regulation difficulties in adolescents using the difficulties in emotion regulation scale. *Assessment*, 17(1), 138–149. <https://doi.org/10.1177/1073191109349579>.
- Noventini, A., Palladino, B. E., & Menesini, E. (2021). Adolescents' stress reactions in response to COVID-19 pandemic at the peak of the outbreak in Italy. *Clinical Psychological Science*, 9(3), 507–514. <https://doi.org/10.1177/2167702621995761>.
- Nolen-Hoeksema, S., Larson, J., & Grayson, C. (1999). Explaining the gender difference in depressive symptoms. *Journal of Personality and Social Psychology*, 77(5), 1061–1072. <https://doi.org/10.1037/0022-3514.77.5.1061>.
- Oelkers, L., Vogel, M., Kalenda, A., Surup, H. C., Körner, A., Kratzsch, J., & Kiess, W. (2020). Socioeconomic status is related to pubertal development in a German cohort. *Hormone Research in Paediatrics*, 93(9–10), 548–557. <https://doi.org/10.1159/000513787>.
- Patel, P. K., Leatham, L. D., Currin, D. L., & Karlsgodt, K. H. (2021). Adolescent neurodevelopment and vulnerability to psychosis. *Biological Psychiatry*, 89(2), 184–193. <https://doi.org/10.1016/j.biopsych.2020.06.028>.
- Pitzer, M. (2019). The development of monoaminergic neurotransmitter systems in childhood and adolescence. *International Journal of Developmental Neuroscience*, 74(1), 49–55. <https://doi.org/10.1016/j.ijdevneu.2019.02.002>.
- Raio, C. M., Oredoru, T. A., Palazzolo, L., Shurick, A. A., & Phelps, E. A. (2013). Cognitive emotion regulation fails the stress test. *Proceedings of the National Academy of Sciences of the United States of America*, 110(37), 15139–15144. <https://doi.org/10.1073/pnas.1305706110>.
- Raio, C. M., & Phelps, E. A. (2015). The influence of acute stress on the regulation of conditioned fear. *Stress Resilience*, 1, 134–146. <https://doi.org/10.1016/j.jynstr.2014.11.004>.
- Scott, B. G., Lapré, G. E., Marsee, M. A., & Weems, C. F. (2014). Aggressive behavior and its associations with posttraumatic stress and academic achievement following a natural disaster. *Journal of Clinical Child & Adolescent Psychology*, 43(1), 43–50. <https://doi.org/10.1080/15374416.2013.807733>.
- Seaton, E., & Carter, R. (2019). Perceptions of pubertal timing and discrimination among African American and Caribbean black girls. *Child Development*, 90. <https://doi.org/10.1111/cdev.13221>.
- Semple, B. D., Blomgren, K., Gimlin, K., Ferriero, D. M., & Noble-Haesuslein, L. J. (2013). Brain development in rodents and humans: Identifying benchmarks of maturation and vulnerability to injury across species. *Progress in Neurobiology*, 106–107, 1–16. <https://doi.org/10.1016/j.pneurobio.2013.04.001>.
- Sowell, E. R., Thompson, P. M., & Toga, A. W. (2004). Mapping changes in the human cortex throughout the span of life. *The Neuroscientist*, 10(4), 372–392. <https://doi.org/10.1177/1073858404263960>.
- Stoet, G. (2010). PsyToolkit: A software package for programming psychological experiments using Linux. *Behavior Research Methods*, 42(4), 1096–1104. <https://doi.org/10.3758/BRM.42.4.1096>.
- Stoet, G. (2017). PsyToolkit: A novel web-based method for running online questionnaires and reaction-time experiments. *Teaching of Psychology*, 44(1), 24–31. <https://doi.org/10.1177/0098628316677643>.
- Thayer, J. F., & Lane, R. D. (2000). A model of neurovisceral integration in emotion regulation and dysregulation. *Arousal in Anxiety*, 61(3), 201–216. [https://doi.org/10.1016/S0165-0327\(00\)00338-4](https://doi.org/10.1016/S0165-0327(00)00338-4).
- Wahlstrom, D., Collins, P., White, T., & Luciana, M. (2010). Developmental changes in dopamine neurotransmission in adolescence: Behavioral implications and issues in assessment. *Brain and Cognition*, 72(1), 146–159. <https://doi.org/10.1016/j.bandc.2009.10.013>.
- Weinberg, A., & Klonsky, E. D. (2009). Measurement of emotion dysregulation in adolescents. *Psychological Assessment*, 21(4), 616–621. <https://doi.org/10.1037/a0016669>.

- Yang, S., & Tseng, K. Y. (2021). Maturation of corticolimbic functional connectivity during sensitive periods of brain development. *Current topics in behavioral neurosciences* (pp. 1–17). Berlin Heidelberg: Springer. https://doi.org/10.1007/7854_2021_239.
- Zhang, C., Ye, M., Fu, Y., Yang, M., Luo, F., Yuan, J., & Tao, Q. (2020). The psychological impact of the COVID-19 pandemic on teenagers in China. *Journal of Adolescent Health, 67* (6), 747–755. <https://doi.org/10.1016/j.jadohealth.2020.08.026>.
- Zhou, S.-J., Zhang, L.-G., Wang, L.-L., Guo, Z.-C., Wang, J.-Q., Chen, J.-C., Liu, M., Chen, X., & Chen, J.-X. (2020). Prevalence and socio-demographic correlates of psychological health problems in Chinese adolescents during the outbreak of COVID-19. *European Child & Adolescent Psychiatry, 29*(6), 749–758. <https://doi.org/10.1007/s00787-020-01541-4>.
- Zimmer-Gembeck, M. J., & Skinner, E. A. (2011). Review: The development of coping across childhood and adolescence: An integrative review and critique of research. *International Journal of Behavioral Development, 35*(1), 1–17. <https://doi.org/10.1177/0165025410384923>.
- Zubovics, E. A., Fiáth, R., Rádosi, A., Pászthy, B., Réthelyi, J. M., Ulbert, I., & Bunford, N. (2021). Neural and self-reported reward responsiveness are associated with dispositional affectivity and emotion dysregulation in adolescents with evidence for convergent and incremental validity. *Psychophysiology, 58*(2) Article e13723. <https://doi.org/10.1111/psyp.13723>.