



# Longitudinal analyses of depression, anxiety, and suicidal ideation highlight greater prevalence in the northern Dutch population during the COVID-19 lockdowns

Anil P.S. Ori<sup>a,b</sup>, Martijn Wieling<sup>c</sup>, Lifelines Corona Research Initiative, Hanna M. van Loo<sup>a,\*</sup>

<sup>a</sup> University of Groningen, University Medical Center Groningen, Department of Psychiatry, Groningen, the Netherlands

<sup>b</sup> University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen, the Netherlands

<sup>c</sup> University of Groningen, Department of Information Science, Groningen, the Netherlands

## ARTICLE INFO

### Keywords:

Major depressive disorder  
Generalized anxiety disorder  
Suicidal ideation  
COVID-19  
Lifelines biobank  
Generalized additive models

## ABSTRACT

**Background:** The pandemic of the coronavirus disease 2019 (COVID-19) has led to an increased burden on mental health.

**Aims:** To investigate the development of major depressive disorder (MDD), generalized anxiety disorder (GAD), and suicidal ideation in the Netherlands during the first fifteen months of the pandemic and three nation-wide lockdowns.

**Method:** Participants of the Lifelines Cohort Study – a Dutch population-based sample – reported current symptoms of MDD and GAD, including suicidal ideation, according to DSM-IV criteria. Between March 2020 and June 2021, 36,106 participants (aged 18–96) filled out a total of 629,811 questionnaires across 23 time points. Trajectories over time were estimated using generalized additive models and analyzed in relation to age, sex, and lifetime history of MDD/GAD.

**Results:** We found non-linear trajectories for MDD and GAD with a higher number of symptoms and prevalence rates during periods of lockdown. The point prevalence of MDD and GAD peaked during the third hard lockdown at 2.88 % (95 % CI: 2.71 %–3.06 %) and 2.92 % (95 % CI: 2.76 %–3.08 %), respectively, in March 2021. Women, younger adults, and participants with a history of MDD/GAD reported significantly more symptoms. For suicidal ideation, we found a significant linear increase over time in younger participants. For example, 20-year-old participants reported 4.14× more suicidal ideation at the end of June 2021 compared to the start of the pandemic (4.64 % (CI: 3.09 %–6.96 %) versus 1.12 % (CI: 0.76 %–1.66 %)).

**Limitations:** Our findings should be interpreted in relation to the societal context of the Netherlands and the public health response of the Dutch government during the pandemic, which may be different in other regions in the world.

**Conclusions:** Our study showed greater prevalence of MDD and GAD during COVID-19 lockdowns and a continuing increase in suicidal thoughts among young adults suggesting that the pandemic and government enacted restrictions impacted mental health in the population. Our findings provide actionable insights on mental health in the population during the pandemic, which can guide policy makers and clinical care during future lockdowns and epi/pandemics.

## 1. Introduction

The pandemic of the coronavirus disease 2019 (COVID-19) had a major impact on societies and led to increases in major depressive disorder (MDD), generalized anxiety disorder (GAD), and suicidality across the world (COVID-19 Mental Disorders Collaborators, 2021; Dubé et al.,

2021a; Robinson et al., 2022a). These conditions are severe and disabling and represent major contributors to the global burden of disease and mortality (Naghavi and Global Burden of Disease Self-Harm Collaborators, 2019). How the prevalence of MDD/GAD and their symptoms in the population changed over time during the COVID-19 pandemic, and which groups are most at risk, especially during

\* Corresponding author at: Department of Psychiatry, University Medical Center Groningen, PO Box 30.001 (CC44), 9700 RB Groningen, the Netherlands.  
E-mail address: [h.van.loo@umcg.nl](mailto:h.van.loo@umcg.nl) (H.M. van Loo).

<https://doi.org/10.1016/j.jad.2022.11.040>

Received 23 August 2022; Received in revised form 2 November 2022; Accepted 18 November 2022

Available online 22 November 2022

0165-0327/© 2022 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

periods of lockdown, remains unclear.

During the first months of the pandemic, overall mental health problems showed a small but significant increase compared to pre-pandemic measures (Penninx et al., 2022). Depressive and anxiety symptoms in particular showed significant increases (Fancourt et al., 2021; Kwong et al., 2021; Prati and Mancini, 2021; Vindegaard and Benros, 2020) that remained elevated for more months for depression (Robinson et al., 2022b). There is however substantial variability in mental health change across studies as well as among subgroups (Penninx et al., 2022; Robinson et al., 2022b). Furthermore, only a handful of longitudinal population-based studies have been conducted that measured MDD/GAD prevalence using psychiatric diagnostic criteria which limits our understanding of how the pandemic and subsequent government-enacted lockdowns impacted clinical outcomes (Demakos, 2021; Penninx et al., 2022). These longitudinal studies were also limited in their estimation of non-linear trajectories. Determining the exact non-linear trajectories of the prevalence of MDD/GAD and their symptoms in the population during the pandemic is thus warranted, as we expect that prevalence rates are dynamic and may increase during lockdowns but may decrease again afterwards. Taking these nonlinear patterns in the development of MDD/GAD and their symptoms over time into account will thus give more fine-grained insight in the potential risk of lockdowns on mental health than estimating average prevalence rates across a certain pre-specified time period. Furthermore, few studies investigated suicidal ideation during the COVID-19 pandemic in a large sample, with no longitudinal study of the general population conducted so far (Farooq et al., 2021), despite that suicidal ideation is strongly associated with depression and anxiety (Cai et al., 2021; Wiebenga et al., 2021), and a major public health concern (Turecki and Brent, 2016).

Here, we investigate the development of (symptoms of) MDD, GAD, and suicidal ideation during the COVID-19 pandemic in Lifelines, a large population-based cohort in the North of the Netherlands. Between April 2020 and July 2021, MDD, GAD and suicidal ideation were repeatedly assessed using DSM-IV criteria in >76,000 participants, which represents the largest longitudinal cohort on pandemic-related impact on mental health in the population (Intyre et al., 2021a). During this period, the Dutch government enacted three nationwide lockdowns, each defined by specific societal restrictions and distancing measures as part of the public health response (National Institute for Public Health and the Environment, 2022). Using detailed self-reported longitudinal data, we first estimated the prevalence of (symptoms of) MDD, GAD, and suicidal ideation across the first fifteen months of the pandemic and three nation-wide lockdown periods using statistical models that allow for non-linear relationships. We then investigated differences in prevalence by age, sex, and lifetime history of MDD/GAD, which are known risk factors of MDD and GAD (Salk et al., 2017; van Loo et al., 2021; Wittchen et al., 1994).

Based on previous findings of increased symptoms of depression and anxiety across the first months of the pandemic and an overall reported increase in MDD, GAD, and suicidality, we hypothesize an increase in prevalence of (symptoms of) MDD, GAD, and suicidal ideation, particularly during periods of lockdown. We furthermore hypothesize a higher prevalence in groups with known risk for developing these symptoms and disorders, i.e. in women, younger adults, and subjects with a lifetime history of MDD/GAD.

## 2. Methods

Full details on cohort information, digital questionnaires, and our analytical strategy can be found in the supplemental materials.

### 2.1. The Lifelines COVID-19 Cohort

Lifelines is a multi-disciplinary prospective population-based cohort study examining in a unique three-generation design the health and health-related behaviors of 167,729 persons living in the North of the

Netherlands. It employs a broad range of investigative procedures in assessing the biomedical, socio-demographic, behavioral, physical and psychological factors which contribute to the health and disease of the general population, with a special focus on multi-morbidity and complex genetics.

In March 2020, the Lifelines Corona Research Initiative was initiated to monitor the physical and mental health of residents in the three Northern provinces of the Netherlands during the COVID-19 pandemic through detailed digital questionnaires (Intyre et al., 2021b). The Lifelines COVID-19 cohort is embedded in Lifelines, a large multi-generational prospective population-based study and biobank with extensive information collected on health, lifestyle and sociodemographic data (Scholtens et al., 2015; Stolk et al., 2008). All participants provided written informed consent. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures involving human subjects were approved by the Medical Ethics Committee of the University Medical Center Groningen, The Netherlands (2007/152).

### 2.2. COVID-19 questionnaires and sample selection

In March 2020, the first digital questionnaire was sent to all 140,145 adult Lifelines participants with an e-mail address on file (Intyre et al., 2021b). Follow-up questionnaires were initially sent on a weekly (questionnaires 1 to 6, Q1-Q6) and later on a biweekly and monthly basis (Q7-Q23). Up to July 2021, 23 questionnaires have been sent out with 76,376 study participants filling in at least one questionnaire (Fig. S1). To minimize the impact of participation bias, i.e. participants with MDD and GAD were less likely to participate in the next questionnaire (see supplementary methods), we selected participants aged 18 years and older who filled out at least one questionnaire in Q1-Q3 and at least one questionnaire in Q21-Q23 to conduct our primary statistical analyses (N = 36,106).

### 2.3. Outcome measures

Current symptoms of MDD and GAD reflecting the DSM-IV criteria (American Psychiatric Association et al., 2000) were assessed using a digital self-report version of the Mini-International Neuropsychiatric Interview (MINI) (Sheehan et al., 1998), which has also been implemented in earlier assessments in Lifelines (van Loo et al., 2021). All items had a binary response option (yes/no) (Table S1, Fig. S2). The first questionnaires (Q1-Q6) assessed current symptoms during the past seven days (as Q1-Q6 were sent out weekly), while the later questionnaires (Q7-Q23) assessed during the past 14 days (as Q7-Q23 were sent out biweekly or monthly). We extracted five outcomes for our analyses. We calculated sum scores for depressive (range 0–9) and anxiety symptoms (range 0–7) that represented a count of the number of MINI items assessing DSM-IV-TR symptom criteria a participant reported. In addition, we used reported symptom expression to determine if a participant met the criteria for MDD and GAD diagnosis according to the DSM-IV-TR, which we coded as a binary outcome measure. Finally, we included reported suicidal ideation as a fifth outcome measure. Suicidal ideation was assessed as present if a participant reported to have considered hurting themselves, wished they were dead, or had suicidal thoughts in the past seven/fourteen days. More information on outcome measures is described in the supplemental methods.

### 2.4. Predictors

We used four predictors in our analyses: time, age, sex assigned at birth, and lifetime history of MDD/GAD (see supplementary methods). Lifetime history of MDD and GAD were determined using an online assessment that is based on the Composite International Diagnostic

Interview (Bot et al., 2017).

## 2.5. Missing data and imputation

To handle missing data, we performed a single dataset imputation using a chained equation regression framework implemented in R-package mice\_v3.13 (van Buuren and Groothuis-Oudshoorn, 2011). We imputed missing values if a participant filled out at least part of that questionnaire. Missingness within filled out questionnaires was overall limited (see supplemental methods). The missing data was imputed using information from other time points within the Lifelines COVID-19 study and from previous assessment waves in Lifelines.

## 2.6. Statistical analyses

Generalized additive models (GAMs) were used to assess the population prevalence of MDD, GAD, and suicidality over time and their association with age, sex, and lifetime history of MDD/GAD. GAMs are regression models that can identify nonlinear patterns in longitudinal data (Wieling, 2018; Wood, 2017). We modeled the prevalence of each of the five MDD/GAD outcomes as a (potentially) non-linear function of time and tested if there were significant interaction effects of time with age, sex, and lifetime history of MDD/GAD. Each outcome and predictor were fitted using a separate model. All analyses were performed in R\_v4.0.3 using the packages mgcv\_1.8.33 (Wood, 2017) and itsadug\_2.4 (van Rij et al., 2015). Multiple testing correction was implemented by Bonferroni correction ( $\alpha = 0.0025$ ).

## 2.7. Sensitivity analyses

We conducted three sensitivity analyses (Fig. S3). First, we implemented all GAMs without random effects in the full cohort ( $N = 76,376$ ) and compared the output with that obtained from our analyses on the main sample of 36,106 subjects. This allowed us to assess the impact of participant dropout on our findings. Second, because individuals were repeatedly assessed over time, we used a random intercept and linear random slope to account for the nested structure of the data within individuals and families. As including random effects for the full cohort was not possible due to computational constraints (see supplementary methods), we conducted the analysis on a subset of 5000 participants (randomly drawn from the 36,106 subjects). While the GAMs with inclusion of random effects estimated lower prevalence (as only fixed effects were returned and the random effects were set to zero), it did allow us to evaluate how individual- and family-specific variation impact the observed effect of predictors and the trajectories over time in a random subset of our sample (see supplementary information). Third, as our main sample of 5000 subjects included a low number of cases for rare phenotypes such as suicidal ideation, particularly for younger ages, we also performed a third sensitivity analysis in the youngest 5000 participants using GAMs with random effects. This analysis included all participants from 18 to 45 years old with at least one assessment in Q1-Q3 and in Q21-Q23.

## 3. Results

### 3.1. Sample description

Our selected sample consisted of 36,106 study participants who completed a total of 629,811 questionnaires with at least 1 questionnaire in Q1-Q3 and at least one questionnaire in Q21-Q23 (Tables 1 and S2). Participants had an average age of 57.4 years ( $SD = 11.9$ ) and filled out a median number of 20 questionnaires. Women (61.9 %) participated more often than men. An average of 1.9 % and 2.3 % of participants met the DSM-IV criteria for current MDD and GAD, respectively, during at least one assessment during the pandemic.

From these participants, a subsample of 5000 subjects was randomly

**Table 1**

Demographics and characteristics of the Lifelines COVID-19 study. Shown are the number of participants and their characteristics of our main analysis sample (36,106 participants). Number of MDD/GAD and suicidal ideation cases are presented as the number of participants who met the DSM-IV criteria for at least one questionnaire. The MDD/GAD prevalence and average symptom scores are presented by the mean and spread of their per-questionnaire average based on the imputed data. Demographics and characteristics of the full cohort and subsamples can be found in Table S2.

	Main analysis sample
Number of participants	36,106
Age	57.4 ( $SD = 11.9$ )
18–30 years	875 (2.4 %)
31–67 years	27,394 (75.9 %)
>68 years	7837 (21.7 %)
Female (%)	22,339 (61.9 %)
Total questionnaires	629,811
Median questionnaire/person (IQR 25 %–75 %)	20 (15–22)
Lifetime MDD (%)	7917 (21.9 %)
Lifetime GAD (%)	2997 (8.3 %)
Number of MDD cases	3675
Average MDD prevalence	1.9 %
Average MDD symptom score	0.50 ( $SD = 1.14$ )
Number of GAD cases	4625
Average GAD prevalence	2.3 %
Average GAD symptom score	0.63 ( $SD = 1.29$ )
Number of suicidal ideation cases	2007
Average suicidal ideation prevalence	0.71 %

drawn to perform sensitivity analyses to assess how individual- and family-specific variation impacts our analyses. The subsample was similar in terms of median number of questionnaires filled out, sex, age distribution, and internalizing disorder distribution to the original sample of 36,106 Lifelines participants (Table S2). Table S2 also shows the characteristics of the full sample and the subsample of the 5000 youngest study participants that we used for sensitivity analyses.

The COVID-19 pandemic and subsequent lockdowns in the Netherlands.

During different phases of the pandemic, the Dutch government enacted a total of three nationwide lockdowns, each defined by specific measures and characteristics (Fig. 1). Using data collected between March 2020 and June 2021, we next estimated the longitudinal trajectories of (symptoms of) MDD/GAD and suicidal ideation across the three lockdown periods.


### 3.2. Longitudinal trajectories of depression, anxiety, and suicidal ideation

#### 3.2.1. Depressive symptoms

We found a significant non-linear trajectory for depressive symptoms over time (Fig. 2, Table S3). Depressive symptoms were high at the start of the pandemic and declined as the first targeted lockdown progressed. Participants reported a lower number of symptoms during mid-summer, which then increased again in August 2020. Symptoms plateaued during the second partial lockdown in November and increased again during the hard lockdown after December 2020. During this third lockdown, reported symptoms reached their peak mid-March 2021 and declined again as the lockdown ended. Comparing the end of April 2021, when restrictions of the hard third lockdown started to be lifted, with the end of April 2020, participants reported more depressive symptoms a year later (0.54 compared to 0.46). The non-linear symptom trajectories were similar in sensitivity analyses in the full cohort as well as after taking into account individual- and family-specific effects (Fig. S4 and Tables S7, S11).

#### 3.2.2. Major depressive disorder

We found a significant non-linear trajectory for the point prevalence of MDD (Fig. 2, Table S3). The prevalence of MDD was relatively stable




**Fig. 1.** Timeline of the COVID-19 pandemic and subsequent government measures in the North of the Netherlands. Shown are the total number of daily reported infections (black bar graphs) and average change in mobility (red line) in the three Northern provinces in the Netherland across three nation-wide lockdowns over time. The number of daily infections were downloaded from the website of the National Institute for Public Health and the Environment (RIVM). The change in mobility was downloaded from Apple Mobility Trends Reports. The change in mobility is expressed as the percentage change compared to a baseline on January 13th, 2020. The time period of the three enacted nation-wide lockdowns are visualized by the gray vertical rectangles. The first known case of a SARS-CoV-2 infection was reported in the Netherlands on February 27 and the first reported death on March 06, 2020. The first lockdown, from March 12–May 31, 2020, to reduce spreading of the virus was defined by targeted measures to restrict social interaction such as closing of public spaces, bars and restaurants, and work from home recommendations. The targeted lockdown reduced mobility in society by 80 % but was not a hard lockdown (de Haas et al., 2020). After the subsequent summer, a new partial lockdown was announced that started on October 14 and lasted until December 14, 2020. During this period of partial lockdown, bars and restaurants were closed and limitations on social gatherings and house visits were recommended. The partial lockdown transitioned into a hard lockdown that lasted for four months which included mandatory closure of all non-essential stores and public space, closure of schools, and an evening curfew, among other measures. The national vaccination program started on January 6, 2021. Easing of restrictions of the hard lockdown were introduced on April 21 and set in motion starting April 28, 2021. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

during the months of 2020 and increased rapidly at the start of the third hard lockdown. We observed a peak in MDD prevalence of 2.88 % (95 % CI: 2.71 %–3.06 %) at the beginning of March 2021, which then declined again the months after. Comparing the end of April 2021 with the end of April 2020, the prevalence of MDD was higher one year later (2.41 % versus 1.78 %;). The trajectory of MDD was similar across sensitivity analyses (Fig. S5 and Tables S7, S11).

### 3.2.3. Anxiety symptoms

Anxiety symptoms showed a significant non-linear trajectory with the highest number of symptoms reported at the immediate start of the pandemic (Fig. 2, Table S3). Reported anxiety symptoms declined as the first lockdown progressed and were lower during mid-summer 2020 when government restrictions were eased. The prevalence of symptoms increased again from August and plateaued during the second lockdown. Anxiety symptoms moderately increased during the third lockdown but did not reach the level of symptoms reported at the start of the pandemic. The prevalence of symptoms declined as the end of the third lockdown approached in May 2021. Comparing the end of April 2021 with the end of April 2020, the number of reported anxiety symptoms one year later was slightly higher (0.65 versus 0.59). Trajectories observed in our sensitivity analyses aligned with our findings from our main analyses (Fig. S6 and Tables S7, S11).



**Fig. 2.** The longitudinal trajectory of MDD and GAD outcomes and suicidal ideation during the COVID-19 pandemic. Shown are the scores of reported depressive and anxiety symptoms and point prevalence of MDD, GAD, and suicidal ideation in the population over time. These trajectories were estimated by GAMS applied to 36,106 Lifelines participants. The x-axis denotes time with the corresponding month and year shown. The gray rectangles highlight the three different nationwide lockdowns in the Netherlands.

### 3.2.4. General anxiety disorder

For GAD, we found a significant non-linear trajectory. The prevalence of GAD was high at the beginning of the pandemic and declined as the first lockdown progressed, reaching its lowest prevalence at the start of July 2020. The months after, the prevalence had a roughly linear increase reaching its peak prevalence of 2.92 % (95 % CI: 2.76 %–3.08 %) at the beginning of March 2021 and declined again toward the end of the third lockdown. Comparing the end of April 2021 with the end of April 2020, the prevalence of GAD was higher one year later (2.71 % versus 2.16 %). The non-linear trajectories were similar in sensitivity analyses in the full cohort as well as after taking into account individual- and family-specific effects (Figs. S5 and S7 and Tables S2, S6, S10). The GAD trajectory did show a more smoothed pattern in the selected subsample of 5,000 participants (Fig. S5), which was likely due to lower statistical power. In the subsample of only younger participants, we observed a more similar trajectory as that of our main analysis.

### 3.2.5. Suicidal ideation

We observed a significant linear increase in the prevalence of suicidal ideation (Fig. 2 and Table S3). At the beginning of April 2020, the prevalence of suicidal ideation was 0.59 % (95 % CI: 0.56 %–0.62 %) which increased to 0.95 % (95 % CI: 0.89 %–1.01 %) during mid-June 2021. This represents a 1.61× increase in reported suicidal ideation in the population. Sensitivity analyses in the full cohort yielded similar results. After taking into account individual-specific and family-specific variation the trajectory of prevalence was flat over time, indicating that most individuals in the population did not experience an increase in suicidal ideation and that specific individuals or subgroups may be more at risk (Fig. S8 and Tables S3, S7, S11).

## 3.3. Longitudinal trajectories across age, sex and history of MDD/GAD

As (symptoms of) MDD and GAD are known to be more prevalent in younger adults, women, and individuals with a previous diagnosis, we next investigated how the observed longitudinal trajectories differed by age, sex, and lifetime history of MDD/GAD in our sample.

### 3.3.1. Longitudinal trajectories by age

Younger participants reported significantly more depressive and anxiety symptoms, as well as a higher prevalence of MDD and GAD than older participants across all time points (Table S4, Figs. S9–S12). The

relative risk of young subjects compared to older subjects did not change across time for these four outcomes nor did they in our sensitivity analyses (Tables S4, S8, S12).

### 3.3.2. Suicidal ideation


At the start of the pandemic, younger participants also reported significantly more suicidal thoughts than older participants (e.g. 1.12 % versus 0.52 % in 20- versus 60-year old participants). Moreover, younger participants also reported a steeper increase in suicidal ideations over time as indicated by a significant interaction effect between time and age (Fig. 3 and Table S4). By mid-June 2021, 20-year-old participants had 4.14× more reports of suicidal ideation than in March 2020 (4.64 % (CI: 3.09 %–6.96 %) versus 1.12 % (CI: 0.76 %–1.66 %)), whereas this increase was lower or absent for older participants. This interaction effect between time and age was also significant in our sensitivity analyses of the youngest 5000 study participants, but not in our randomly selected subsample (Tables S8 and S12, Figs. S14 and S15). As suicidal ideation is a rarer phenotype, including more young adults increased our statistical power to detect such an effect.

### 3.3.3. Longitudinal trajectories by sex


As expected, the prevalence of MDD and GAD and their symptom scores were on average significantly higher in women than in men (similarly to findings in (van Loo et al., 2021)), but the development of these outcomes over time did not differ between the sexes (Figs. S16–S25 and Tables S5, S9, S13). The prevalence and development of suicidal thoughts across time did not differ between women and men, which we also observed in our sensitivity analyses.

### 3.3.4. Longitudinal trajectories by lifetime history of MDD/GAD

Study participants with a lifetime history of MDD/GAD reported significantly more symptoms and a higher prevalence of MDD/GAD and suicidal ideation during the pandemic than participants without a previous diagnosis (Fig. 4, Table S6). We found a significant difference in trajectories over time between participants with and without a history for MDD or GAD for depressive or anxiety symptom scores, respectively, while not for the other three outcomes (Table S6). For both depressive and anxiety symptom scores, the difference between participants with and without a lifetime diagnosis was greatest during periods out of lockdown when reported symptoms were lowest in the general population (Fig. 4). Sensitivity analyses in the full cohort yielded similar



**Fig. 3.** The trajectory of suicidal ideation by time and age during the COVID-19 pandemic. Shown are the results of interaction between time and age on reported suicidal ideation in our main sample of 36,106 participants (age 18–45). The left panels show the trajectory of suicidal ideation for specific ages over time. The right panels show the trajectory of specific time points across age. The legends at the top of the graph denoted colour coding of groups. The gray rectangles highlight the three different nationwide lockdowns in the Netherlands.



**Fig. 4.** The trajectory of depressive and anxiety symptom scores between study participants with and without a history of MDD/GAD during the COVID-19 pandemic. Shown are the results of interaction between time and a lifetime history of MDD/GAD on symptom scores based on GAM analyses in our main analysis sample. The left panels show the trajectory of the mean score of reported depressive and anxiety symptoms between participants with (green) and without (blue) a history of MDD/GAD. The right panels show corresponding difference plots that visualize the trajectory of the difference between the two groups. The gray rectangles highlight the three different nationwide lockdowns in the Netherlands. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

findings, as did correcting for individual- and family-specific variation (Figs. S26–S35 and Tables S6, S10, S14).

#### 4. Discussion

To investigate the impact of the COVID-19 pandemic on mental health, we described the development of current (symptoms of) MDD, GAD and suicidal thoughts in a longitudinal sample from the Northern Dutch general population who were followed for more than a year during the COVID-19 pandemic across three nation-wide lockdowns.

##### 4.1. Longitudinal trajectories of depression, anxiety, and suicidal ideation

In general, we observed a greater prevalence of (symptoms of) MDD and GAD in the population during periods of lockdown and a declining prevalence during periods of eased or no restrictions, which suggests that the pandemic and government measures enacted to restrict spreading of the coronavirus indeed impacted the mental health of the general population. We observed a high prevalence of symptoms of MDD and GAD at the start of the pandemic followed by a rapid decline in the months after, which is in line with observations from other countries (Batterham et al., 2021; Fancourt et al., 2021; Kwong et al., 2021; Robinson et al., 2022b). A new finding of our study is that the prevalence of symptoms of MDD and GAD increased again during later lockdowns alongside a high prevalence of MDD and GAD during the third lockdown which peaked in March 2021. The third lockdown in the Netherlands, unlike the first and second lockdown, was a hard lockdown and characterized by stricter measures, including an evening curfew.

Interestingly, we observed a plateauing of reported symptoms during the second lockdown, which was a shorter partial lockdown, where schools, sports, libraries and museums, unlike during the third lockdown, stayed open. Together our results suggest that lockdown measures impacted mental health in the population, although we submit that this is an observational study of multifactorial psychiatric conditions.

##### 4.2. Longitudinal trajectories across age, sex and history of MDD/GAD

Women, younger adults, and subjects with pre-existing mental health conditions, were more at risk for developing MDD and GAD symptoms and disorders, which is in line with observations from previous studies as well (Batterham et al., 2021; Fancourt et al., 2021; Kwong et al., 2021). However, apart from a difference in intercept, we observed no differences in the development of MDD/GAD over time across sex or age, meaning that the relative risk of MDD/GAD between men and women, or between older and younger participants did not change across time. We did find a difference in the trajectory of reported MDD/GAD symptoms between participants with a history of MDD/GAD compared to participants without a history, but this difference was not found for MDD and GAD. Participants with a history of MDD/GAD reported increased symptom severity during the end of the first lockdown and months after, which is in contrast to a smaller Dutch study that found no difference at the start of the pandemic (Pan et al., 2021) and highlights the importance of continued longitudinal measurements in large cohorts.

In contrast to the non-linear trajectories of (symptoms of) MDD and GAD, the prevalence of suicidal ideation showed an increasing linear

trend among young adults while we did not observe this effect in the overall population. There were no sex differences in prevalence rates of suicidality, but subjects with a lifetime history of MDD reported more suicidal ideation. A meta-analysis of suicidality during the pandemic found increased rates of suicidal ideation and behaviors with a higher incidence in younger individuals as well (Dubé et al., 2021b). As only 14 % of studies on suicidality during the pandemic included children or young people (John et al., 2020), there is an urgent need for more epidemiological studies, like the Lifelines COVID-19 study, that includes these groups to investigate what factors are driving the observed increase. As psychological distress, low perceived social support, and loneliness are known factors that increase suicidality in adolescents and young adults (McClelland et al., 2020; Yildiz, 2020), government measures to reduce the spread of the virus by social distancing measures likely contributed to the increased prevalence. Adolescents and young adults were furthermore significantly impacted by unemployment at the start of the pandemic (Dutch Central Bureau of Statistics, 2021). As unemployment increases risk of suicide (Kawohl and Nordt, 2020), our observation of increased suicidal thoughts may translate to suicide attempts and mortality (Dubé et al., 2021b). Indeed, the Dutch Suicide Prevention Center reported an increase in completed suicides among young adults in January and February of 2021 compared to previous years (Dutch National Suicide Prevention, 2021). These alarming findings warrant for alertness in psychiatric care services and urge governments to consider the long-term impact of pandemic measurements on young people.

#### 4.3. Strengths and limitations

A key strength of our study is the high-quality and high-resolution longitudinal data collected using a validated structured diagnostic interview throughout the first fifteen months of the pandemic spanning three nationwide lockdowns. This allowed for the application of sophisticated nonlinear statistical models to investigate the development of MDD, GAD and suicidality across time. However, our findings should be interpreted considering several limitations. First, due to the large computational resources required to run GAMs with random effects, we were only able to account for individual- and family-specific variation in a subsample of our cohort. While this subsample was similar to our larger sample in its main characteristics, we cannot exclude that our analyses may have missed important insights due to limited statistical power, especially for more rare phenotypes. We did show that analyses on the youngest study participants is worthwhile to prioritize as a significant proportion of the variation in MDD/GAD outcomes lies in younger adults, which increases statistical power to identify group differences. Second, we fitted each outcome in a separate model and thus could only compare the prevalence within a single trajectory and not between trajectories of different outcomes. As depression and anxiety are known to have comorbidity, how changes in trajectories relate among outcomes is an important question to investigate in future research. Third, we did not have information on the prevalence of MDD, GAD and suicidality in the year before the start of the pandemic and thus could not account for that, nor for seasonal effects. Fourth, we assessed current symptoms of GAD within the past seven and fourteen days. We therefore did not assess GAD according to the DSM-IV criteria that requires symptoms to be present for at least six months. Fifth, we did not account for corona infection status of study participants. As the number of participants who reported to have tested positive for the coronavirus during the study was relatively low (11.2 %), we expect infection status and its accompanying symptoms to have minimal impact on our findings, if any at all, given that the observed trajectories in prevalence of MDD/GAD and suicidal ideation also do not follow the number of reported infections over time (Fig. 1). Sixth, Lifelines participants are more often female, middle aged, married, and Dutch native compared to the population in the North of the Netherlands (Klijs et al., 2015). We therefore cannot exclude that there are population sub-groups who may

be at greater risk of declining mental health that are under-represented in the Lifelines COVID 19 study. Finally, our findings should be interpreted within the societal context of the study. The Netherlands had three lockdown periods with different characteristics and is furthermore on average a rich country with a social welfare system.

## 5. Conclusion

In summary, we investigated the development of (symptoms of) MDD and GAD and suicidal thoughts in the Northern Dutch population during the COVID-19 pandemic and observed higher prevalence during periods of lockdown, in particular the third hard lockdown. We furthermore found an alarming linear increase in suicidal thoughts among young adults that warrants for alertness in psychiatric care services. Further studies are needed to investigate mechanisms underlying these rising prevalence rates. Our findings provide important insights into the impact of the pandemic on the mental health of the population, which can help guide policy makers and clinical care during future lockdowns and *epi*/pandemics.

### Funding support

HvL was supported by a NARSAD Young Investigator Grant from the Brain & Behaviour Research Foundation and a VENI grant from the Talent Programme of the Netherlands Organization of Scientific Research (NWO-ZonMW 09150161810021). The funding organizations had no impact on study design, data collection, analysis, or interpretation, or decision to submit the manuscript.

### CRediT authorship contribution statement

All authors contributed to the design of the study. The Lifelines Corona Research Initiative, AO and HvL were involved in data collection. AO, HvL and MW designed the study. AO performed the statistical analyses. AO drafted the manuscript; all other authors interpreted results and provided feedback on analyses and drafts of the manuscript.

### Conflict of Interest

The authors report no conflicts of interest.

### Data availability

All data were obtained from the Lifelines biobank under project application OV20.00021. Data can be obtained by submitting a request to the Lifelines Research Office ([research@lifelines.nl](mailto:research@lifelines.nl)). The data are not publicly available because they contain information that could compromise the study participants' privacy and consent.

### Acknowledgements

The authors wholeheartedly thank the study participants for their time and invaluable contribution to the Lifelines Biobank.

The Lifelines Biobank initiative has been made possible by funding from the Dutch Ministry of Health, Welfare and Sport, the Dutch Ministry of Economic Affairs, the University Medical Center Groningen (UMCG the Netherlands), the University of Groningen, the Northern Provinces of the Netherlands, FES (Fonds Economische Structuurversterking), SNN (Samenwerkingsverband Noord Nederland) and REP (Ruimtelijk Economisch Programma).

We acknowledge funding for the Lifelines Corona Research project from the University of Groningen and the University Medical Centre Groningen. The authors wish to acknowledge the efforts of the Lifelines Corona Research Initiative and the following initiatives participants:

HM Boezen<sup>†</sup> (1), Jochen O. Mierau (2,3), H. Lude Franke (4), Jackie Dekens (4,6), Patrick Deelen (4), Pauline Lanting (4), Judith M. Vonk

(1), Ilja Nolte (1), Anil P.S. Ori (4,5), Annique Claringbould (4), Floranne Boulogne (4), Marjolein X.L. Dijkema (4), Henry H. Wiersma (4), Robert Warmerdam (4), Soesma A. Jankipersadzing (4). Irene van Blokland (4,7).

1) Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

2) Faculty of Economics and Business, University of Groningen, Groningen, The Netherlands.

3) Aletta Jacobs School of Public Health, Groningen, The Netherlands.

4) Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

5) Department of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

6) Center of Development and Innovation, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

7) Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

We thank the UMCG Genomics Coordination Center, the UG Center for Information Technology and their sponsors BBMRI-NL & TarGet for storage and computing infrastructure.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jad.2022.11.040>.

## References

- American Psychiatric Association, American Psychiatric Association Staff, American Psychiatric Association, 2000. *Task Force on DSM-IV. In: Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR. Amer Psychiatric Pub Incorporated.*
- Batterham, P.J., Calear, A.L., McCallum, S.M., Morse, A.R., Banfield, M., Farrer, L.M., Gulliver, A., Cherbuin, N., Rodney Harris, R.M., Shou, Y., Dawel, A., 2021. Trajectories of depression and anxiety symptoms during the COVID-19 pandemic in a representative Australian adult cohort. *Med. J. Aust.* <https://doi.org/10.5694/mja.251043>.
- Bot, M., Middeldorp, C.M., de Geus, E.J.C., Lau, H.M., Sinke, M., van Nieuwenhuizen, B., Smit, J.H., Boomsma, D.I., Penninx, B.W.J.H., 2017. Validity of LIDAS (Lifetime Depression Assessment Self-report): a self-report online assessment of lifetime major depressive disorder. *Psychol. Med.* 47, 279–289.
- Cai, H., Jin, Y., Liu, S., Zhang, Q., Zhang, L., Cheung, T., Balbuena, L., Xiang, Y.-T., 2021. Prevalence of suicidal ideation and planning in patients with major depressive disorder: a meta-analysis of observation studies. *J. Affect. Disord.* 293, 148–158.
- COVID-19 Mental Disorders Collaborators, 2021. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet* 398, 1700–1712.
- de Haas, M., de Haas, M., Faber, R., Hamersma, M., 2020. How COVID-19 and the Dutch “intelligent lockdown” change activities, work and travel behaviour: evidence from longitudinal data in the Netherlands. *Transportation Research Interdisciplinary Perspectives.* <https://doi.org/10.1016/j.trip.2020.100150>.
- Demakos, P., 2021. Importance of population-based longitudinal studies to understanding the impact of COVID-19. *J. Epidemiol. Community Health* 75, 815–816.
- Dubé, J.P., Smith, M.M., Sherry, S.B., Hewitt, P.L., Stewart, S.H., 2021a. Suicide behaviors during the COVID-19 pandemic: a meta-analysis of 54 studies. *Psychiatry Res.* 301, 113998.
- Dubé, J.P., Smith, M.M., Sherry, S.B., Hewitt, P.L., Stewart, S.H., 2021b. Suicide behaviors during the COVID-19 pandemic: a meta-analysis of 54 studies. *Psychiatry Res.* 301, 113998.
- Dutch Central Bureau of Statistics, 2021. Youth labour participation recovered partially in Q3–Q4 2020 [WWW Document]. Statistics Netherlands. <https://www.cbs.nl/en-gb/news/2021/09/youth-labour-participation-recovered-partially-in-q3-q4-2020>.
- Dutch National Suicide Prevention, 2021. Geen toename in totaal aantal suicides, extra aandacht voor jongeren noodzakelijk [WWW Document]. URL <https://www.113.nl/actueel/geen-toename-totaal-aantal-suicides-extra-aandacht-voor-jongeren-noodzakelijk>.
- Fancourt, D., Steptoe, A., Bu, F., 2021. Trajectories of anxiety and depressive symptoms during enforced isolation due to COVID-19 in England: a longitudinal observational study. *Lancet Psychiatry* 8, 141–149.
- Farooq, S., Tunmore, J., Wajid Ali, M., Ayub, M., 2021. Suicide, self-harm and suicidal ideation during COVID-19: a systematic review. *Psychiatry Res.* 306, 114228.
- Intyre, K.M., Mc Intyre, K., Lanting, P., Deelen, P., Wiersma, H.H., Vonk, J.M., Ori, A.P. S., Jankipersadzing, S.A., Warmerdam, R., van Blokland, I., Boulogne, F., Dijkema, M.X.L., Herkert, J.C., Claringbould, A., Bakker, O., Lopera Maya, E.A., Bültmann, U., Zernakova, A., Reijneveld, S.A., Zijlstra, E., Swertz, M.A., Brouwer, S., van Ooijen, R., Angelini, V., Dekker, L.H., Sijtsma, A., Scherjon, S.A., Wijmenga, C., Dekens, J.A.M., Mierau, J., Marike Boezen, H., Franke, L., 2021a. Lifelines COVID-19 cohort: investigating COVID-19 infection and its health and societal impacts in a Dutch population-based cohort. *BMJ Open.* <https://doi.org/10.1136/bmjopen-2020-044474>.
- Intyre, K.M., Mc Intyre, K., Lanting, P., Deelen, P., Wiersma, H.H., Vonk, J.M., Ori, A.P. S., Jankipersadzing, S.A., Warmerdam, R., van Blokland, I., Boulogne, F., Dijkema, M.X.L., Herkert, J.C., Claringbould, A., Bakker, O., Lopera Maya, E.A., Bültmann, U., Zernakova, A., Reijneveld, S.A., Zijlstra, E., Swertz, M.A., Brouwer, S., van Ooijen, R., Angelini, V., Dekker, L.H., Sijtsma, A., Scherjon, S.A., Wijmenga, C., Dekens, J.A.M., Mierau, J., Marike Boezen, H., Franke, L., 2021b. Lifelines COVID-19 cohort: investigating COVID-19 infection and its health and societal impacts in a Dutch population-based cohort. *BMJ Open.* <https://doi.org/10.1136/bmjopen-2020-044474>.
- John, A., Eyles, E., Webb, R.T., Okolie, C., Schmidt, L., Arensman, E., Hawton, K., O'Connor, R.C., Kapur, N., Moran, P., O'Neill, S., McGuiness, L.A., Olorisade, B.K., Dekel, D., Macleod-Hall, C., Cheng, H.-Y., Higgins, J.P.T., Gunnell, D., 2020. The impact of the COVID-19 pandemic on self-harm and suicidal behaviour: update of living systematic review. *F1000Res.* 9, 1097.
- Kawohl, W., Nordt, C., 2020. COVID-19, unemployment, and suicide. *Lancet Psychiatry.* [https://doi.org/10.1016/s2215-0366\(20\)30141-3](https://doi.org/10.1016/s2215-0366(20)30141-3).
- Klijs, B., Scholtens, S., Mandemakers, J.J., Snieder, H., Stolk, R.P., Smidt, N., 2015. Representativeness of the Lifelines Cohort Study. *PLoS One* 10, e0137203.
- Kwong, A.S.F., Pearson, R.M., Adams, M.J., Northstone, K., Tilling, K., Smith, D., Fawcitt-Ritchie, C., Bould, H., Warne, N., Zammit, S., Gunnell, D.J., Moran, P.A., Micali, N., Reichenberg, A., Hickman, M., Rai, D., Haworth, S., Campbell, A., Altschul, D., Flaig, R., McIntosh, A.M., Lawlor, D.A., Porteous, D., Timpson, N.J., 2021. Mental health before and during the COVID-19 pandemic in two longitudinal UK population cohorts. *Br. J. Psychiatry* 218 (6), 334–343. <https://doi.org/10.1192/bjp.2020.242>.
- McClelland, H., Evans, J.J., Nowland, R., Ferguson, E., O'Connor, R.C., 2020. Loneliness as a predictor of suicidal ideation and behaviour: a systematic review and meta-analysis of prospective studies. *J. Affect. Disord.* 274, 880–896.
- Naghavi, M., Global Burden of Disease Self-Harm Collaborators, 2019. Global, regional, and national burden of suicide mortality 1990 to 2016: systematic analysis for the Global Burden of Disease Study 2016. *BMJ* 364, 194.
- National Institute for Public Health and the Environment, 2022. Timeline of the COVID-19 pandemic in the Netherlands [WWW Document]. <https://www.rivm.nl/gedragsonderzoek/tijdlijn-maatregelen-covid> accessed 10.30.22.
- Pan, K.-Y., Kok, A.A.L., Eikelenboom, M., Horsfall, M., Jörg, F., Luteijn, R.A., Rhebergen, D., van Oppen, P., Giltay, E.J., Penninx, B.W.J.H., 2021. The mental health impact of the COVID-19 pandemic on people with and without depressive, anxiety, or obsessive-compulsive disorders: a longitudinal study of three Dutch case-control cohorts. *Lancet Psychiatry* 8, 121–129.
- Penninx, B.W.J.H., Benros, M.E., Klein, R.S., Vinkers, C.H., 2022. How COVID-19 shaped mental health: from infection to pandemic effects. *Nat. Med.* 28, 2027–2037.
- Prati, G., Mancini, A.D., 2021. The psychological impact of COVID-19 pandemic lockdowns: a review and meta-analysis of longitudinal studies and natural experiments. *Psychol. Med.* 51, 201–211.
- Robinson, E., Sutin, A.R., Daly, M., Jones, A., 2022a. A systematic review and meta-analysis of longitudinal cohort studies comparing mental health before versus during the COVID-19 pandemic in 2020. *J. Affect. Disord.* 296, 567–576.
- Robinson, E., Sutin, A.R., Daly, M., Jones, A., 2022b. A systematic review and meta-analysis of longitudinal cohort studies comparing mental health before versus during the COVID-19 pandemic in 2020. *J. Affect. Disord.* 296, 567–576.
- Salk, R.H., Hyde, J.S., Abramson, L.Y., 2017. Gender differences in depression in representative national samples: meta-analyses of diagnoses and symptoms. *Psychol. Bull.* 143, 783–822.
- Scholtens, S., Smidt, N., Swertz, M.A., Bakker, S.J.L., Dotinga, A., Vonk, J.M., van Dijk, F., van Zon, S.K.R., Wijmenga, C., Wolfenbuttel, B.H.R., Stolk, R.P., 2015. Cohort profile: Lifelines, a three-generation cohort study and biobank. *Int. J. Epidemiol.* 44, 1172–1180.
- Sheehan, D.V., Lecrubier, Y., Sheehan, K.H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R., Dunbar, G.C., 1998. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J. Clin. Psychiatry* 59 (Suppl. 20), 22–33 quiz 34–57.
- Stolk, R.P., Rosmalen, J.G.M., Postma, D.S., de Boer, R.A., Navis, G., Slaets, J.P.J., Ormel, J., Wolfenbuttel, B.H.R., 2008. Universal risk factors for multifactorial diseases: Lifelines: a three-generation population-based study. *Eur. J. Epidemiol.* 23, 67–74.
- Turecki, G., Brent, D.A., 2016. Suicide and suicidal behaviour. *Lancet* 387, 1227–1239.
- van Buuren, S., Groothuis-Oudshoorn, K., 2011. mice: multivariate imputation by chained equations in R. *J. Stat. Softw.* 45, 1–67.
- van Loo, H.M., Beijers, L., Wieling, M., de Jong, T.R., Schoevers, R.A., Kendler, K.S., 2021. Prevalence of internalizing disorders, symptoms, and traits across age using advanced nonlinear models. *Psychol. Med.* 1–10.
- van Rij, J., Wieling, M., Baayen, R.H., van Rijn, H., 2015. itsadug: interpreting time series and autocorrelated data using GAMMS.
- Vindegard, N., Benros, M.E., 2020. COVID-19 pandemic and mental health consequences: systematic review of the current evidence. *Brain Behav. Immun.* 89, 531–542.
- Wiebenga, J.X.M., Dickhoff, J., Mérelle, S.Y.M., Eikelenboom, M., Heering, H.D., Gilissen, R., van Oppen, P., Penninx, B.W.J.H., 2021. Prevalence, course, and determinants of suicide ideation and attempts in patients with a depressive and/or anxiety disorder: a review of NESDA findings. *J. Affect. Disord.* 283, 267–277.



Wieling, M., 2018. Analyzing dynamic phonetic data using generalized additive mixed modeling: a tutorial focusing on articulatory differences between L1 and L2 speakers of English. *J. Phon.* <https://doi.org/10.1016/j.wocn.2018.03.002>.

Wittchen, H.U., Zhao, S., Kessler, R.C., Eaton, W.W., 1994. DSM-III-R generalized anxiety disorder in the National Comorbidity Survey. *Arch. Gen. Psychiatry* 51, 355–364.

Wood, S.N., 2017. *Generalized Additive Models: An Introduction With R*, Second edition. CRC Press.

Yıldız, M., 2020. Stressful life events and adolescent suicidality: an investigation of the mediating mechanisms. *J. Adolesc.* 82, 32–40.